Carcinoembryonic antigen peptide 1

Cat. No.: HY-P0277 CAS No.: 168635-85-6 Molecular Formula: $C_{43}H_{69}N_{11}O_{14}$

Molecular Weight: 964.07

Tyr-Leu-Ser-Gly-Ala-Asn-Leu-Asn-Leu Sequence:


Sequence Shortening: YLSGANLNL Target: Others Pathway: Others

Sealed storage, away from moisture and light Storage:

> Powder -80°C 2 years -20°C 1 year

* In solvent: -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture

and light)



Product Data Sheet

SOLVENT & SOLUBILITY

In Vitro

H₂O: 12.5 mg/mL (12.97 mM; Need ultrasonic)

Preparing Stock Solutions	Solvent Mass Concentration	1 mg	5 mg	10 mg
	1 mM	1.0373 mL	5.1863 mL	10.3727 mL
	5 mM	0.2075 mL	1.0373 mL	2.0745 mL
	10 mM	0.1037 mL	0.5186 mL	1.0373 mL

Please refer to the solubility information to select the appropriate solvent.

BIOLOGICAL ACTIVITY

Description	Carcinoembryonic antigen peptide 1 (CAP1) is a tumor marker in lung cancer.
In Vitro	Carcinoembryonic Antigen (CEA) as tumor marker in lung cancer. The use of CEA as a prognostic and predictive marker. CEA is a glycoprotein involved in cell adhesion, and is normally produced during fetal development but the production stops before birth. CEA is a glycosyl phosphatidyl inositol (GPI)-cell surface anchored glycoprotein whose specialized sialofucosylated glycoforms serve as functional colon carcinoma L-selectin and E-selectin ligands, which may be critical to the metastatic dissemination of colon carcinoma cells ^[1] . MCE has not independently confirmed the accuracy of these methods. They are for reference only.

REFERENCES

1]. Grunnet M, et al. Carcinoembryo	onic antigen (CEA) as tumor marker in lung cancer. Lung Ca	ncer. 2012 May;76(2):138-43.
Ca	aution: Product has not been fully validated for medi	cal applications. For research use only.
Tel	el: 609-228-6898 Fax: 609-228-5909 Address: 1 Deer Park Dr, Suite Q, Monmout	E-mail: tech@MedChemExpress.com th Junction, NJ 08852, USA

Page 2 of 2 www.MedChemExpress.com