

Lys- γ 3-MSH(human) TFA

Cat. No.:	HY-P1210A
Molecular Formula:	C ₁₃₀ H ₁₉₄ F ₃ N ₄₅ O ₄₁ S
Molecular Weight:	3132.27
Sequence Shortening:	KYVMGHFRWDRFGRRNSSSSGSSGAGQ
Target:	Others
Pathway:	Others
Storage:	Please store the product under the recommended conditions in the COA.

BIOLOGICAL ACTIVITY

Description

Lys- γ 3-MSH(human) TFA is a melanocortin peptide derived from the C-terminal of the fragment of pro-opiomelanocortin (POMC). Lys- γ 3-MSH(human) TFA potentiates the steroidogenic response of the rat adrenal to adrenocorticotrophin (ACTH). Lys- γ 3-MSH(human) TFA is a potent stimulator of lipolysis with an apparent EC₅₀ of 3.56 nM. Lys- γ 3-MSH(human) TFA can activate hormone sensitive lipase (HSL) and Perilipin A resulting in lipolysis^{[1][2]}

REFERENCES

- [1]. Harmer SC, Pepper DJ, Cooke K, Bennett HP, Bicknell AB. Evidence of a possible role for Lys-gamma3-MSH in the regulation of adipocyte function. *J Endocrinol.* 2008;196(1):149-158.
- [2]. Bicknell KA, et al. Lys-gamma3-MSH: a global regulator of hormone sensitive lipase activity?. *Mol Cell Endocrinol.* 2009;300(1-2):71-76.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA