

Apraglutide TFA

Cat. No.:	HY-P1714A
Molecular Formula:	$C_{172}H_{263}N_{43}O_{52} \cdot C_2HF_3O_2$
Molecular Weight:	3879.27
Sequence Shortening:	HGDGSFSDEX FTILDLLAAR DFINWLIQTK ITD-NH2
Target:	GLP Receptor
Pathway:	GPCR/G Protein
Storage:	Sealed storage, away from moisture Powder -80°C 2 years -20°C 1 year * In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture)

BIOLOGICAL ACTIVITY

Description	Apraglutide TFA (FE 203799 TFA), a synthetic 33-amino-acid peptide and a long-acting GLP-2 analogue, enhances adaptation and linear intestinal growth in a neonatal piglet model of short bowel syndrome with total resection of the ileum ^[1] .								
In Vivo	<p>Apraglutide (FE 203799; 5 mg/kg/dose, subcutaneously, twice on days 0 and 4 postsurgery) treated piglets are healthy, have significant lower fecal fat and energy losses and exhibit intestinal lengthening, greater small-intestinal weight, longer villus height, and greater crypt depth on day 7^[1].</p> <p>MCE has not independently confirmed the accuracy of these methods. They are for reference only.</p> <table border="1"> <tr> <td>Animal Model:</td> <td>NewbornDuroc piglets, 2-5 days old and weighing between 2-2.6 kg^[1].</td> </tr> <tr> <td>Dosage:</td> <td>5 mg/kg/dose.</td> </tr> <tr> <td>Administration:</td> <td>Subcutaneously, twice on days 0 and 4 postsurgery.</td> </tr> <tr> <td>Result:</td> <td>On day 7, treated piglets were healthy, had significant lower fecal fat and energy losses and exhibited intestinal lengthening, greater small-intestinal weight, longer villus height, and greater crypt depth.</td> </tr> </table>	Animal Model:	NewbornDuroc piglets, 2-5 days old and weighing between 2-2.6 kg ^[1] .	Dosage:	5 mg/kg/dose.	Administration:	Subcutaneously, twice on days 0 and 4 postsurgery.	Result:	On day 7, treated piglets were healthy, had significant lower fecal fat and energy losses and exhibited intestinal lengthening, greater small-intestinal weight, longer villus height, and greater crypt depth.
Animal Model:	NewbornDuroc piglets, 2-5 days old and weighing between 2-2.6 kg ^[1] .								
Dosage:	5 mg/kg/dose.								
Administration:	Subcutaneously, twice on days 0 and 4 postsurgery.								
Result:	On day 7, treated piglets were healthy, had significant lower fecal fat and energy losses and exhibited intestinal lengthening, greater small-intestinal weight, longer villus height, and greater crypt depth.								

REFERENCES

[1]. Slim GM, et al. Novel Long-Acting GLP-2 Analogue, FE 203799 (Apraglutide), Enhances Adaptation and Linear Intestinal Growth in a Neonatal Piglet Model of Short Bowel Syndrome with Total Resection of the Ileum. JPEN J Parenter Enteral Nutr. 2019 Jan 6.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA