

Product Data Sheet

Animal-Free NAP-2/CXCL7 Protein, Mouse (His)

Cat. No.:	HY-P700174AF
Synonyms:	C-X-C motif chemokine 7; PBP; LDGF; MDGF; CTAP-III; PPBP; NAP-2; CXCL7
Species:	Mouse
Source:	E. coli
Accession:	Q9EQI5 (K40-Y113)
Gene ID:	57349
Molecular Weight:	Approximately 8.93 kDa

PROPERTIES	
AA Sequence	KSDGMDPYIE LRCRCTNTIS GIPFNSISLV NVYRPGVHCA DVEVIATLKN GQKTCLDPNA PGVKRIVMKI LEGY
Biological Activity	Measure by its ability to chemoattract BaF3 cells transfected with human CXCR2. The ED ₅₀ for this effect is <1 μ g/mL.
Appearance	Lyophilized powder.
Formulation	Lyophilized from a solution containing 1X PBS, pH 7.4.
Endotoxin Level	<0.1 EU per 1 μ g of the protein by the LAL method.
Reconsititution	It is not recommended to reconstitute to a concentration less than 100 $\mu\text{g}/\text{mL}$ in ddH_2O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US;may vary elsewhere.

DESCRIPTION

 Background
 The NAP-2/CXCL7 protein is a member of the intercrine alpha (chemokine CxC) family. This classification highlights its

 affiliation with a group of chemokines involved in intercellular communication and immune responses. As part of the

 intercrine alpha family, NAP-2/CXCL7 likely plays a role in modulating inflammatory processes and cellular interactions.

 Further exploration is necessary to uncover the specific functions and implications of this protein within the broader context

 of the chemokine CxC family, shedding light on its significance in mediating immune responses without reliance on animal

 derived sources.

Caution: Product has not been fully validated for medical applications. For research use only.

 Tel: 609-228-6898
 Fax: 609-228-5909
 E-mail: tech@MedChemExpress.com

 Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA