

## Animal-Free BMP-4 Protein, Pig (His)

<b>Cat. No.:</b>	HY-P700230AF
<b>Synonyms:</b>	BMP-2B; BMP-4; Bone morphogenetic protein 4; DVR4
<b>Species:</b>	Pig
<b>Source:</b>	E. coli
<b>Accession:</b>	A0A6I9IR93 (R43-C224)
<b>Gene ID:</b>	100113425
<b>Molecular Weight:</b>	Approximately 21.65 kDa

### PROPERTIES

<b>AA Sequence</b>	<pre> M R E K Q P N Y G L   A I E V T H L H Q T   R T H Q G Q H V R I   S R S L P Q G S G D W A Q L R P L L V T   F G H D G R G H A L   T R R R R A K R S P   K H H P Q R A R K K N K N C R R H S L Y   V D F S D V G W N D   W I V A P P G Y Q A   F Y C H G D C P F P L A D H L N S T N H   A I V Q T L V N S V   N S S I P K A C C V   P T E L S A I S M L Y L D E Y D K V V L   K N Y Q E M V V E G   C G C           </pre>
<b>Appearance</b>	Lyophilized powder
<b>Formulation</b>	Lyophilized from a solution containing 20 mM sodium citrate, 0.2 M NaCl pH 3.5.
<b>Endotoxin Level</b>	<0.1 EU per 1 µg of the protein by the LAL method.
<b>Reconstitution</b>	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH <sub>2</sub> O.
<b>Storage &amp; Stability</b>	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
<b>Shipping</b>	Room temperature in continental US; may vary elsewhere.

### DESCRIPTION

<b>Background</b>	<p>Bone Morphogenetic Protein 4 (BMP-4) is a ligand protein with pleiotropic, belongs to TGFβ family. BMP-4 involves in the vasculature circulation and can activate receptors on vascular cells<sup>[1]</sup>.</p> <p>BMP-4/TGFβ signaling can be terminated by inhibitory SMADs including SMAD6 and SMAD7, which are activated and induced by BMP signaling and switch off BMP signaling via multiple mechanisms<sup>[4]</sup>.</p> <p>BMP-4 is widely found in different animals, while the sequence in human is highly similar to Rat (96.81%), and mouse (97.54%).</p> <p>BMP-4 is expressed by endothelial cells (ECs) in response to hypoxia and promotes vascular SMC proliferation. Therefore it inhibits the proliferation of smooth muscle cells (SMCs) isolated from the proximal pulmonary artery while induces</p>
-------------------	--

---

proliferation of SMCs isolated from distal pulmonary arteries<sup>[5]</sup>.  
BMP-4 appears to be a marker and driver of vascular calcification, particularly in atherosclerosis<sup>[6]</sup>.  
BMP-4 induces angiogenesis, endothelial cells (ECs) proliferation, and migration<sup>[7]</sup>.  
BMP-4 is differentially expressed in calcified atherosclerotic plaques<sup>[8]</sup>, serves as the linkers between atherosclerotic vascular calcification with mechanisms of normal bone formation<sup>[9]</sup>.  
BMP-4 increases plaque formation via their pro-inflammatory and pro-atherogenic effects, promoting oxidative stress, endothelial dysfunction and osteogenic differentiation<sup>[3]</sup>.

---

## REFERENCES

---

- [1]. Yang P, et al. The role of bone morphogenetic protein signaling in vascular calcification. *Bone*. 2020 Dec;141:115542.
- [2]. Miyazawa K, et al. Regulation of TGF- $\beta$  Family Signaling by Inhibitory Smads. *Cold Spring Harb Perspect Biol*. 2017 Mar 1;9(3):a022095.
- [3]. Herrera B, et al. A rapid and sensitive bioassay for the simultaneous measurement of multiple bone morphogenetic proteins. Identification and quantification of BMP4, BMP6 and BMP9 in bovine and human serum. *BMC Cell Biol*. 2009 Mar 19;10:20.
- [4]. Yang X, et al. Dysfunctional Smad signaling contributes to abnormal smooth muscle cell proliferation in familial pulmonary arterial hypertension. *Circ Res*. 2005 May 27;96(10):1053-63.
- [5]. Scimeca M, et al. Plaque calcification is driven by different mechanisms of mineralization associated with specific cardiovascular risk factors. *Nutr Metab Cardiovasc Dis*. 2019 Dec;29(12):1330-1336.
- [6]. David L, et al. Emerging role of bone morphogenetic proteins in angiogenesis. *Cytokine Growth Factor Rev*. 2009 Jun;20(3):203-12.
- [7]. Dhore CR, et al. Differential expression of bone matrix regulatory proteins in human atherosclerotic plaques. *Arterioscler Thromb Vasc Biol*. 2001 Dec;21(12):1998-2003.
- [8]. Demer LL, et al. Mechanism of calcification in atherosclerosis. *Trends Cardiovasc Med*. 1994 Jan-Feb;4(1):45-9.
- [9]. Boström K, et al. Bone morphogenetic protein expression in human atherosclerotic lesions. *J Clin Invest*. 1993 Apr;91(4):1800-9.
- 

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA