

Animal-Free FGF-2 Protein, Pig (His)

Cat. No.:	HY-P700237AF
Synonyms:	Fibroblast growth factor 2; FGF-2; Basic fibroblast growth factor; bFGF; Heparin-binding growth factor 2; HBGF-2; FGF2; FGFB
Species:	Pig
Source:	E. coli
Accession:	NP_001392443.1
Gene ID:	397643
Molecular Weight:	Approximately 18.1 kDa

PROPERTIES

AA Sequence	A A G S I T T L P A L P E D G G S G A F P P G H F K D P K R L Y C K N G G F F L R I H P D G R V D G V R E K S D P H I K L Q L Q A E E R G V V S I K G V C A N R Y L A M K E D G R L L A S K C V T D E C F F F E R L E S N N Y N T Y R S R K Y S S W Y V A L K R T G Q Y K L G P K T G P G Q K A I L F L P M S A K S
Biological Activity	Measure by its ability to induce proliferation in 3T3 cells. The ED ₅₀ for this effect is <2 ng/mL
Appearance	Lyophilized powder
Formulation	Lyophilized from a solution containing 0.01% sarkosyl in 1X PBS, pH 7.4.
Endotoxin Level	<0.1 EU per 1 µg of the protein by the LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background	FGF-2/bFGF is a member of the fibroblast family and has a high affinity for heparin. FGF-2 plays an important role in tendon to bone healing, cartilage repair, bone repair, and nerve regeneration. FGF-2 specifically binds to tyrosine kinase receptors and activates the FGF/FGFR signaling pathway. Subsequently, FGF-2 influences cell proliferation, differentiation and apoptosis, as well as immune regulation by transducing other classical pathways. For example, FGF-2 regulates the JAK-STAT signaling pathway to regulate cartilage metabolism. FGF-2 also acts as a mitotic promoter to accelerate cell proliferation. Therefore, (1) FGF-2 is an important growth factor in the healing process of ligament/tendon injury. In vitro experiments, low-dose FGF-2 can stimulate the proliferation and differentiation of bone marrow mesenchymal stem cells,
------------	---

and up-regulate the mRNA expression of type I/III collagen and fibronectin. However, high doses of FGF-2 did not stimulate extracellular matrix (ECM) protein proliferation and gene expression. (2) FGF-2 is also an endogenous and intrinsic growth factor in cartilage repair. FGF-2 binds to heparan sulfate proteoglycan and is stored in the ECM of articular cartilage. When cartilage is damaged or degenerated, ECM rapidly releases FGF-2 and activates ERK signaling pathways to promote cartilage regeneration. FGF-2 exhibits a biphasic effect in combination with its specific receptor. FGF-2 combined with FGFR3 promoted the repair of articular cartilage. FGF-2 combined with FGFR1 promoted the degeneration of articular cartilage^[1]. FGF-2 is expressed in granulosa cells and colliculus cells, as well as hepatocellular cancer cells, but not in non-cancerous liver tissues. This reveals the role of FGF-2 in brain tumors, particularly glioblastoma. According to studies, FGF-2 is a known carcinogenic factor in GBM. FGF-2 increases the self-renewal of glioblastoma stem cells and contributes to the growth and vascularization of glioma^[2]. FGF-2 protein is highly conserved in some species, and the similarity rate of human FGF-2 protein sequence to rat, mouse, and bovine was 97.4%, 95.45%, and 98.71%, respectively.

REFERENCES

- [1]. Zhang J, et al. FGF2: a key regulator augmenting tendon-to-bone healing and cartilage repair. *Regen Med.* 2020 Sep;15(9):2129-2142.
- [2]. Jimenez-Pascual A, et al. FGF2: a novel druggable target for glioblastoma? *Expert Opin Ther Targets.* 2020 Apr;24(4):311-318.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA