

Animal-Free GM-CSF Protein, Pig (His)

Cat. No.:	HY-P700239AF
Synonyms:	CSF2
Species:	Pig
Source:	E. coli
Accession:	Q29118 (A18-K144)
Gene ID:	397208
Molecular Weight:	Approximately 15.3 kDa

PROPERTIES

AA Sequence	<p>A P T R P P S P V T R P W Q H V D A I K E A L S L L N N S N D T A A V M N E T V</p> <p>D V V C E M F D P Q E P T C V Q T R L N L Y K Q G L R G S L T R L K S P L T L L</p> <p>A K H Y E Q H C P L T E E T S C E T Q S I T F K S F K D S L N K F L F T I P F D</p> <p>C W G P V K K</p>
Biological Activity	Measure by its ability to induce proliferation in TF-1 cells. The ED ₅₀ for this effect is <3 ng/mL.
Appearance	Lyophilized powder.
Formulation	Lyophilized from a solution containing 1X PBS, pH 7.4.
Endotoxin Level	<0.1 EU per 1 µg of the protein by the LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background	GM-CSF Protein serves as a potent cytokine, orchestrating the growth and differentiation of hematopoietic precursor cells across diverse lineages, including granulocytes, macrophages, eosinophils, and erythrocytes. Structurally, GM-CSF exists as a monomer, and its signaling is mediated through a dodecameric complex. This complex comprises two head-to-head hexamers, each featuring two alpha, two beta, and two ligand subunits, underscoring the intricacy of the receptor assembly and the specificity of GM-CSF in modulating cellular responses.
------------	--

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA