RedChemExpress

Product Data Sheet

protein G protein (C-His)

Cat. No.:	HY-P700321
Synonyms:	Immunoglobulin G-binding protein G; spg; IgG-binding protein G
Species:	Staphylococcus aureus
Source:	E. coli
Accession:	P19909 (L298-E49)
Gene ID:	/
Molecular Weight:	Approximately 27 kDa

PROPERTIES Biological Activity Measured by its binding ability in a functional ELISA. Immobilized Protein G at 2µg/mL (100µL/well) can bind Biotinylated Human IgG1. The ED ₅₀ for this effect is 0.1099 µg/mL. Appearance Lyophilized powder Formulation Lyophilized a 0.2 µm filtered solution of PBS, pH 7.4, 5 % trehalose, 5 % mannitol and 0.01% Tween80. Endotoxin Level -1 EU/µg, determined by LAL method.
Human IgG1. The ED ₅₀ for this effect is 0.1099 μg/mL. Appearance Formulation Lyophilized a 0.2 μm filtered solution of PBS, pH 7.4, 5% trehalose, 5% mannitol and 0.01% Tween80.
Formulation Lyophilized a 0.2 μm filtered solution of PBS, pH 7.4, 5 % trehalose, 5 % mannitol and 0.01% Tween80.
Endotoxin Level <1 EU/μg, determined by LAL method.
ReconsititutionIt is not recommended to reconstitute to a concentration less than 100 μ g/mL in ddH2O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).
Storage & Stability Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping Room temperature in continental US; may vary elsewhere.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898 Fax: 609-228-5909

28-5909 E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA