

## Product Data Sheet

## CD24 Protein-Nanoparticle, Human (HEK293, His)

| Cat. No.:         | HY-P700401   |
|-------------------|--|
| Synonyms:         | CD24; CD24 molecule; CD24 antigen (small cell lung carcinoma cluster 4 antigen); signal<br>transducer CD24; CD24A; FLJ22950; FLJ43543; MGC75043; |
| Species:          | Human  |
| Source:           | HEK293   |
| Accession:        | P25063 (S27-G59)   |
| Gene ID:          | 100133941  |
| Molecular Weight: | 25.7 kDa   |

| PROPERTIES          |  |
|---------------------|--|
| AA Sequence         | SETTTGTSSN SSQSTSNSGL APNPTNATTK AAG   |
| Appearance          | Lyophilized powder.  |
| Formulation         | Lyophilized from a 0.2 $\mu m$ filtered solution of 20 mM Tris-HCl, 0.5 M NaCl, 6% Trehalose, pH 8.0   |
| Endotoxin Level     | <1 EU/µg, determined by LAL method.  |
| Reconsititution     | It is not recommended to reconstitute to a concentration less than 100 $\mu\text{g}/\text{mL}$ in ddH_2O.  |
| Storage & Stability | Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage. |
| Shipping            | Room temperature in continental US; may vary elsewhere.  |

| DESCRIPTION |   |
|-------------|---|
| Background  | The CD24 protein is implicated in playing a pivotal role in the cell differentiation of various cell types, with its signaling potentially triggered by the binding of a lectin-like ligand to CD24 carbohydrates, transduced through the release of second messengers derived from the GPI-anchor. CD24 also functions in modulating B-cell activation responses, promoting AG-dependent proliferation of B-cells, and preventing their terminal differentiation into antibody-forming cells. In association with SIGLEC10, CD24 may contribute to the selective suppression of the immune response to danger-associated molecular patterns (DAMPs) such as HMGB1, HSP70, and HSP90, suggesting a role in immune regulation. Additionally, CD24 is |
|             | implicated in the control of autoimmunity. It exhibits cell-type-specific interactions with LYN and FGR, binding to FRG in a<br>small cell cancer line, LYN in the erythroleukemia cell line K-562, and in Burkitt's lymphoma, highlighting its diverse<br>involvement in cellular processes and immune regulation.   |

## Caution: Product has not been fully validated for medical applications. For research use only.

 Tel: 609-228-6898
 Fax: 609-228-5909
 E-mail: tech@MedChemExpress.com

 Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA