

Tau-F/MAPT Protein, Human (HEK293, His)

Cat. No.:	HY-P700421
Synonyms:	MAPT; TAU; MSTD; PPND; DDPAC; MAPTL; MTBT1; MTBT2; FTDP-17; microtubule-associated protein tau; microtubule-associated protein tau; PHF-tau; paired helical filament-tau; neurofibrillary tangle protein; microtubule-associated protein tau, isoform 4; G protein beta1/gamma2 subunit-interacting factor 1
Species:	Human
Source:	HEK293
Accession:	P10636-8 (M1-L441)
Gene ID:	4137
Molecular Weight:	49.5 kDa

PROPERTIES

AA Sequence	<pre> MAEPRQEFEV MEDHAGTYGL GDRKDQGGYT MHQDQEGDTD AGLKEsplQT PTEdGSEEPG SETSDAKSTP TAEDVTAPLV DEGAPGKQAA AQPHTeIPEG TTAEEAGIGD TPSLEDEAAG HVTQARMVSK SKDGTGSDDK KAKGADGKTK IATPRGAAPP GQKQGQANATR IPAKTPPAPK TPPSSGEPK SGDRSGYSSP GSPGTPGSRs RTPSLPTPPT REPKKVAVVR TPPKSPSSAK SRLQTAPVPM PDLKNVKSki GSTENLKHQP GGGKVQIINK KLDLSNVQSK CGSKDNIKHV PGGGSVQIVY KPVDLSKVTS KCGSLGNIHH KPGGGQVEVK SEKLDFKDRV QSKIGSLDNI THVPGGGNKK IETHKLTFRE NAKAKTDHGA EIVYKSPVVS GDTSPRHLsN VSSTGSIDMV DSPQLATLAD EVSASLAKQG L </pre>
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 µm filtered solution of 20 mM Tris-HCl, 0.5 M NaCl, 6% Trehalose, pH 8.0
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background

Microtubule-associated protein tau is a microtubule-associated protein that is found in large quantities in neurons of the central nervous system (CNS). MAPT promotes microtubule assembly and stability and may be involved in the establishment and maintenance of neuronal polarity. MAPT can bind axon microtubules at the C end and neurotic membrane components at the N end, acting as a connexion between the two. Defects in MAPT can lead to neurological diseases such as Alzheimer's and Parkinson's. Overexpression of MAPT is associated with a poor prognosis of prostate cancer. Tau protein S262/S356 can be phosphorylated by AMPK-related kinase^{[1][2][3][4]}.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA