

Tau/MAPT Protein, Mouse (HEK293, His)

Cat. No.:	HY-P700422
Synonyms:	MAPT; TAU; MSTD; PPND; DDPAC; MAPTL; MTBT1; MTBT2; FTDP-17; microtubule-associated protein tau; microtubule-associated protein tau; PHF-tau; paired helical filament-tau; neurofibrillary tangle protein; microtubule-associated protein tau, isoform 4; G protein beta1/gamma2 subunit-interacting factor 1
Species:	Mouse
Source:	HEK293
Accession:	P10637 (A2-L733)
Gene ID:	17762
Molecular Weight:	78.9 kDa

PROPERTIES

AA Sequence

```

ADPRQEFDTM    EDHAGDYTL    LDQDEGDMDHG  LKESPPQPPA
DDGAEPEGSE    TSDAKSTPTA  EDVTAPLVDE   RAPDKQAAAQ
PHTEIPEGIT    AEEAGIGDTP  NQEDQAAGHV   TQGRREGQAP
DLGTSDWTRQ    QVSSMSGAPL  LPQGLREATC   QPSGTRPEDI
EKSHPASELL    RRGPPQKEGW  GQDRLGSEEE   VDEDLTVDES
SQDSPPSQAS    LTPGRAAPQA  GSGSVCGETA   SVPGLPTEGS
VPLPADFFSK    VSAETQASQP  EPGTGPMEE    GHEAAPEFTF
HVEIKASTPK    EQDLEGATVV  GVPGEEQKAQ   TQGPSVGKGT
KEASLQEPG    KQPAAGLPGR  PVS RVPQLKA  RVASKDRTGN
DEKKAKTSTP    SCAKAPSHRP  CLSPTRPTLG   SSDPLIKPSS
PAVSPEPATS    PKHVSSVTPR  NGSPGTKQMK   LKGADGKTGA
KIATPRGAAS    PAQKGTSNAT  RIPAKTTPSP   KTPPGSGEPP
KSGERSGYSS    PGSPGTPGSR  SRTPSLPTPP   TREPKKVAVV
RTPPKSPSAS    KSRLQTAPVP  MPDLKNVRSK   IGSTENLKHQ
PGGGKVQIIN    KKLDSLNVQS  KCGSKDNIKH   VPGGGSVQIV
YKPVDSLKVT    SKCGSLGNIH  HKPGGGQVEV   KSEKLDKFDR
VQSKIGSLDN    ITHVPGGGNK  KIETHKLTFR   ENAKAKTDHG
AEIVYKSPVV    SGDTSRHL    NVSSTGSIDM   VDSPQLATLA
DEVSASLAKQ    GL
  
```

Appearance

Lyophilized powder.

Formulation

Lyophilized from a 0.2 µm filtered solution of 20 mM Tris-HCl, 0.5 M NaCl, 6% Trehalose, pH 8.0

Endotoxin Level

<1 EU/µg, determined by LAL method.

Reconstitution

It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH₂O.

Storage & Stability

Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is

recommended to freeze aliquots at -20°C or -80°C for extended storage.

Shipping

Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background

Microtubule-associated protein tau (MAPT) is a key player in the promotion of microtubule assembly and stability, suggesting its potential involvement in the establishment and maintenance of neuronal polarity. Its C-terminus binds to axonal microtubules, while the N-terminus interacts with neural plasma membrane components, indicating tau's role as a crucial linker protein bridging these cellular structures. The predetermined axonal polarity is dictated by tau's localization within the neuronal cell, specifically in the domain defined by the centrosome. Short isoforms of MAPT contribute to the plasticity of the cytoskeleton, whereas longer isoforms may preferentially play a role in its stabilization. MAPT engages in diverse interactions with various proteins such as MARK1, MARK2, MARK3, MARK4, SQSTM1, PSMC2, FKBP4, CSNK1D, SGK1, EPM2A, PIN1, LRRK2, and LRP1, participating in processes ranging from ubiquitination to dephosphorylation and endocytosis. These intricate interactions highlight MAPT's multifaceted involvement in cellular dynamics.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA