

SSTR2 Protein-VLP, Human (HEK293, His)

Cat. No.:	HY-P700430
Synonyms:	SSTR2; somatostatin receptor 2; somatostatin receptor type 2; SS2R; SRIF-1; SS-2-R
Species:	Human
Source:	HEK293
Accession:	P30874 (M1-I369)
Gene ID:	6752
Molecular Weight:	42.5 kDa

PROPERTIES

AA Sequence	MDMADEPLNG SHTWLSIPFD LNGSVVSTNT SNQTEPYDDL T SNAVLTFIY FVVCIIGLCG NTLVIYVILR YAKMKTITNI YILNLAIADE LFMLGLPFLA MQVALVHWP GKAICRVVMT VDGINQFTSI FCLTVMSIDR YLAVVHPK AKWRRPRTAK MITMAVWGV LLVILPIMY AGLRSNQWGR SSCTINWPGE SGAWYTGFI YTFILGFLVP LTIICLCYLF I I I KVKSSGI RVGSSKRKKS EKKVTRMVS VVAVFIFCWL PFYIFNVSSV SMAISPTPAL KGMFDFVVVL TYANSCANPI LYAFLSDNFK KSFQNVLC KVS GTDDGER SDSKQDKSRL NETTETQRTL LNGDLQTSI
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, 6% Trehalose, pH 7.4.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA