

CD20/MS4A1 Protein-VLP, Dog (HEK293, His)

Cat. No.:	HY-P700444
Synonyms:	Ms4a1; Cd20; Ly-44; Ms4a2; B-lymphocyte antigen CD20; B-cell differentiation antigen Ly-44; Lymphocyte antigen 44; CD antigen
Species:	Dog
Source:	HEK293
Accession:	Q3C2E2 (M1-P297)
Gene ID:	485430
Molecular Weight:	34.4 kDa

PROPERTIES

AA Sequence	<pre> MTT PRNSMSG TLPVDPMKSP TAMYPVQKII PKRMPSVVG P TQNFFMRESK TLGAVQIMNG LFHIALGSL L MIHTDVCAP I CITMWYPLWG GIMFIIISGSL LAAADKNPRK SLVKGKMIM N SLSLFAAISG IIFLIMDIFN ITISHFFKME NLNLIKAPM P YVDIHNCDPA NPSEKNSLSI QYCGSIRSVF LGVFAVMLI F AFFQKLVTAG IVENEWKKLC SKPKSDVVVL LAAEEKKEQ P IETTEEMVEL TEIASQPKKE EDIEIIPVQE EEGELEINFA EPPQE QESS P IENDSIP </pre>
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, 6% Trehalose, pH 7.4.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA