

CD31/PECAM-1 Protein, Cynomolgus (HEK293, His)

| | |
|-------------------|---|
| Cat. No.: | HY-P700675 |
| Synonyms: | PECAM-1; EndoCAM; PECA1; CD31; FLJ34100; FLJ58394; GPIIA' |
| Species: | Cynomolgus |
| Source: | HEK293 |
| Accession: | A0A7N9CHZ3 (H28-G602) |
| Gene ID: | / |
| Molecular Weight: | 95-115 kDa |

PROPERTIES

| | |
|---------------------|--|
| Appearance | Lyophilized powder. |
| Formulation | Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4. Normally 8% trehalose is added as protectant before lyophilization. |
| Endotoxin Level | <1 EU/µg, determined by LAL method. |
| Reconstitution | It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O. |
| Storage & Stability | Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage. |
| Shipping | Room temperature in continental US; may vary elsewhere. |

DESCRIPTION

| | |
|------------|--|
| Background | <p>CD31/PECAM-1 protein, a pivotal cell adhesion molecule, is essential for leukocyte transendothelial migration (TEM) in most inflammatory conditions. The critical role of Tyr-679 in TEM is underscored by its requirement for efficient trafficking of PECAM1 to and from the lateral border recycling compartment (LBRC), crucial for targeting the LBRC membrane around migrating leukocytes. The trans-homophilic interaction potentially contributes to endothelial cell-cell adhesion via cell junctions, while heterophilic interaction with CD177 plays a role in the transendothelial migration of neutrophils. Homophilic ligation of PECAM1 serves a dual purpose, preventing macrophage-mediated phagocytosis of neighboring viable leukocytes by transmitting a detachment signal, while promoting the macrophage-mediated phagocytosis of apoptotic leukocytes by tethering them to phagocytic cells. Notably, PECAM1 modulates bradykinin receptor BDKRB2 activation and regulates bradykinin- and hyperosmotic shock-induced ERK1/2 activation in endothelial cells. Its interaction with various partners, including BDKRB2, GNAQ, PTPN11, FER, and CD177, further elucidates the intricate molecular mechanisms governing its diverse cellular functions, while its trans-homodimerization is crucial for effective cell-cell interaction. The interaction nuances, such as Ca(2+)-dependency and directness, add layers of complexity to the regulatory roles of CD31/PECAM-1.</p> |
|------------|--|

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA