

Fractalkine/CX3CL1 Protein, Cynomolgus (HEK293, His)

Cat. No.:	HY-P700702
Synonyms:	Fractalkine; Neurotactin; Cx3cl1; Cx3c; Fkn; ABCD-3; C3Xkine; CXC3; CXC3C; NTN; NTT; SCYD1
Species:	Cynomolgus
Source:	HEK293
Accession:	EHH60415.1 (Q25-Q341)
Gene ID:	/
Molecular Weight:	50-70 kDa

PROPERTIES

Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.22 μ m filtered solution of PBS, pH 7.4. Normally 8% trehalose is added as protectant before lyophilization.
Endotoxin Level	<1 EU/ μ g, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 μ g/mL in ddH ₂ O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background	Fractalkine/CX3CL1 protein is a member of the intercrine beta (chemokine CC) family, positioning it within a category of chemokines with pivotal roles in intercellular communication and immune responses. As part of the intercrine beta family, Fractalkine/CX3CL1 likely plays a significant role in modulating inflammatory processes and cellular interactions. Further exploration is essential to uncover the specific functions and implications of Fractalkine/CX3CL1 within the broader framework of the chemokine CC family, shedding light on its importance in mediating immune responses and contributing to the intricate network of intercellular signaling.
------------	---

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA