

CD30/TNFRSF8 Protein, Mouse (263a.a, HEK293, His)

Cat. No.:	HY-P70075
Synonyms:	rMuTumor necrosis factor receptor superfamily member 8/TNFRSF8, His; Tumor necrosis factor receptor superfamily member 8; CD30L receptor; Lymphocyte activation antigen CD30; CD30; Tnfrsf8
Species:	Mouse
Source:	HEK293
Accession:	Q60846 (F19-T281)
Gene ID:	21941
Molecular Weight:	54-63 kDa

PROPERTIES

AA Sequence	<pre> F P T D R P L K T T C A G D L S H Y P G E A A R N C C Y Q C P S G L S P T Q P C P R G P A H C R K Q C A P D Y Y V N E D G K C T A C V T C L P G L V E K A P C S G N S P R I C E C Q P G M H C C T P A V N S C A R C K L H C S G E E V V K S P G T A K K D T I C E L P S S G S G P N C S N P G D R K T L T S H A T P Q A M P T L E S P A N D S A R S L L P M R V T N L V Q E D A T E L V K V P E S S S S K A R E P S P D P G N A E K N M T L E L P S P G T L P D I S T S E N S K E P A S T A S T L S L V V D A W T S S R M Q P T S P L S T G T </pre>
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background	CD30/TNFRSF8 Protein, serving as the receptor for TNFSF8/CD30L, plays a critical role in the regulation of cellular growth and the transformation of activated lymphoblasts. Additionally, it is involved in the modulation of gene expression through the activation of NF-kappa-B signaling, contributing to diverse cellular processes. The interaction with key signaling molecules, including TRAF1, TRAF2, TRAF3, and TRAF5, further underscores its significance in mediating intracellular pathways essential for cellular responses and regulatory functions.
-------------------	--

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA