

## **Product** Data Sheet

# PVRIG Protein, Mouse (HEK293, Fc-Avi)

Cat. No.: HY-P700819

Synonyms: C7orf15; CD112R; MGC104322; MGC138297; MGC2463; PVRIG; ALS2CR18; ALS2CR9; LPD; PREL-2;

PREL2; RalGDS/AF-6; RMO1

Species: Mouse Source: **HEK293** 

Accession: A0A1B0GS01 (S35-D165)

Gene ID: 102640920 Molecular Weight: 53-63 kDa

### **PROPERTIES**

Biological Activity	Immobilized Mouse Nectin-2, His Tag at $2\mu g/ml$ ( $100\mu l/well$ ) on the plate. Dose response curve for Mouse PVRIG, hFc Tag with the EC $_{50}$ of $0.61 ug/ml$ determined by ELISA.
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4. Normally 8% trehalose is added as protectant before lyophilization.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconsititution	It is not recommended to reconstitute to a concentration less than 100 $\mu g/mL$ in ddH <sub>2</sub> O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

#### **DESCRIPTION**

### **Background**

Poliovirus receptor related immunoglobulin domain containing (PVRIG), a member of the nectin and nectin-like family, is an immune checkpoint molecule with potential for development. In humans, PVRIG is expressed on T cells (predominantly CD8 <sup>†</sup> T cells) and natural killer (NK) cells, but not on B cells, monocytes or neutrophils. PVRIG binds to a single ligand, poliovirus receptor-related 2 (PVRL2), and exerts an inhibitory effect on cytotoxic lymphocyte activity, likely via an ITIM-like motif in its intracellular domain. PVRIG binds with high affinity to PVRL2 are inhibitory receptors on effector T cells, suppressing cytokine production and cytotoxic activity. PVRIG deficiency or PVRIG blockade can reduce the tumor size and prolong the survival of tumor-bearing mice through inhibiting NK cell and CD8<sup>+</sup> T cell exhaustion. PVRIG blockade enhances natural killer cell killing of PVRL2hiPVRlo acute myeloid leukemia cells<sup>[1][2][3]</sup>.

Page 1 of 2 www.MedChemExpress.com  $\label{lem:caution:Product} \textbf{Caution: Product has not been fully validated for medical applications. For research use only.}$ 

Tel: 609-228-6898 Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA

Page 2 of 2 www.MedChemExpress.com