

Tenascin/Tnc protein, Human (HEK293, His)

Cat. No.:	HY-P700833
Synonyms:	TN; Hexabrachion; Tenascin-C; TN-C; TNC; Hxb
Species:	Human
Source:	HEK293
Accession:	P24821-1 (G23-S621)
Gene ID:	3371
Molecular Weight:	Approximately 53-80 kDa due to the glycosylation.

PROPERTIES

AA Sequence	<pre> G V L K K V I R H K R Q S G V N A T L P E E N Q P V V F N H V Y N I K L P V G S Q C S V D L E S A S G E K D L A P P S E P S E S F Q E H T V D G E N Q I V F T H R I N I P R R A C G C A A A P D V K E L L S R L E E L E N L V S S L R E Q C T A G A G C C L Q P A T G R L D T R P F C S G R G N F S T E G C G C V C E P G W K G P N C S E P E C P G N C H L R G R C I D G Q C I C D D G F T G E D C S Q L A C P S D C N D Q G K C V N G V C I C F E G Y A G A D C S R E I C P V P C S E E H G T C V D G L C V C H D G F A G D D C N K P L C L N N C Y N R G R C V E N E C V C D E G F T G E D C S E L I C P N D C F D R G R C I N G T C Y C E E G F T G E D C G K P T C P H A C H T Q G R C E E G Q C V C D E G F A G V D C S E K R C P A D C H N R G R C V D G R C E C D D G F T G A D C G E L K C P N G C S G H G R C V N G Q C V C D E G Y T G E D C S Q L R C P N D C H S R G R C V E G K C V C E Q G F K G Y D C S D M S C P N D C H Q H G R C V N G M C V C D D G Y T G E D C R D R Q C P R D C S N R G L C V D G Q C V C E D G F T G P D C A E L S C P N D C H G Q G R C V N G Q C V C H E G F M G K D C K E Q R C P S D C H G Q G R C V D G Q C I C H E G F T G L D C G Q H S C P S D C N N L G Q C V S G R C I C N E G Y S G E D C S </pre>
Biological Activity	Measured by the ability of the immobilized protein to block Fibronectin-mediated adhesion of NIH-3T3 mouse embryonic fibroblast cells. Tenascin immobilized at 15 µg/mL, in the presence of 0.1 µg/mL human Fibronectin, will block approximately 72.51% NIH-3T3 cell adhesion (5×10 ⁴ cells/well, 100 µL/well).
Appearance	Lyophilized powder
Formulation	Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.

Shipping

Room temperature in continental US; may vary elsewhere.

DESCRIPTION**Background**

The TNC protein, an extracellular matrix protein, plays a crucial role in guiding migrating neurons and axons during development, contributing to processes such as synaptic plasticity and neuronal regeneration. It facilitates neurite outgrowth from cortical neurons grown on a monolayer of astrocytes, indicative of its involvement in the intricate cellular interactions within the nervous system. Acting as a ligand for integrins alpha-8/beta-1, alpha-9/beta-1, alpha-V/beta-3, and alpha-V/beta-6, TNC establishes molecular connections essential for cellular communication and signaling. In the context of tumors, TNC stimulates angiogenesis by promoting the elongation, migration, and sprouting of endothelial cells. Structurally, TNC exists as a homohexamer, with a potential homotrimer formation in the triple coiled-coil region, further stabilized by disulfide rings at both ends. This versatile protein also interacts with CSPG4, indicating its involvement in diverse cellular and molecular processes across different biological contexts.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA