

TPBG/5T4 Protein, Human (FITC, HEK293, His-Avi)

Cat. No.:	HY-P700840
Synonyms:	M6P1; 5T4AG; WAIF1; TPBG; 5T4
Species:	Human
Source:	HEK293
Accession:	Q13641-1 (S32-S355)
Gene ID:	7162
Molecular Weight:	50-70 kDa

PROPERTIES

Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.22 μ m filtered solution of PBS, pH 7.4. Normally 8% trehalose is added as protectant before lyophilization.
Endotoxin Level	<1 EU/ μ g, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 μ g/mL in ddH ₂ O.
Storage & Stability	Stored at -20°C for 1 year, protect from light. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background	The TPBG/5T4 protein appears to serve as an inhibitor of Wnt/beta-catenin signaling, potentially achieved through indirect interaction with LRP6, thereby impeding Wnt3a-dependent LRP6 internalization. This implies a crucial role for TPBG/5T4 in modulating the intricate Wnt/beta-catenin signaling pathway, exerting regulatory influence on cellular responses associated with this pathway. Further exploration of the specific molecular mechanisms governing the interaction between TPBG/5T4 and LRP6, as well as its impact on Wnt3a-dependent LRP6 internalization, could yield valuable insights into the functional significance of TPBG/5T4 in shaping Wnt/beta-catenin signaling dynamics and its potential implications for various cellular processes.
------------	---

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA