

CD3 epsilon Protein, Cynomolgus (HEK293, Fc)

Cat. No.:	HY-P70090
Synonyms:	rCynT-cell surface glycoprotein CD3 epsilon chain/CD3E, Fc ; CD3 epsilon; CD3e antigen; CD3e antigen, epsilon polypeptide (TiT3 complex); CD3e molecule, epsilon (CD3-TCR complex); CD3e; CD3-epsilon; FLJ18683; T3E; T-cell antigen receptor complex, epsilon subunit of T3; T-cell surface antigen T3/Leu-4 epsilon chain;
Species:	Cynomolgus
Source:	HEK293
Accession:	Q95LI5 (Q22-D117)
Gene ID:	102133065
Molecular Weight:	38-55 kDa

PROPERTIES

AA Sequence	<p>Q D G N E E M G S I T Q T P Y Q V S I S G T T V I L T C S Q H L G S E A Q W Q H</p> <p>N G K N K E D S G D R L F L P E F S E M E Q S G Y Y V C Y P R G S N P E D A S H</p> <p>H L Y L K A R V C E N C M E M D</p>
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 µm filtered solution of 50 mM Tris-HCl, 100 mM Glycine, pH 7.5.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background	<p>The CD3E/CD3 epsilon 1-27 peptide, integral to the TCR-CD3 complex on T-lymphocyte cell surfaces, plays a crucial role in adaptive immune responses. Upon activation by antigen-presenting cells (APCs), the T-cell receptor (TCR) transmits signals across the cell membrane through CD3 chains, including CD3D, CD3E, CD3G, and CD3Z, each containing immunoreceptor tyrosine-based activation motifs (ITAMs) in their cytoplasmic domains. Phosphorylation of these ITAMs by Src family protein tyrosine kinases LCK and FYN, upon TCR engagement, activates downstream signaling pathways. Beyond its role in signal transduction, CD3E is indispensable for correct T-cell development, initiating the assembly of the TCR-CD3 complex by forming heterodimers CD3D/CD3E and CD3G/CD3E. Additionally, CD3E participates in the internalization and cell surface down-regulation of TCR-CD3 complexes through endocytosis sequences in its cytosolic region. The TCR-CD3 complex,</p>
-------------------	--

comprising CD3D/CD3E and CD3G/CD3E heterodimers, preferentially associates with TCRalpha and TCRbeta, forming trimers. This hexamer then interacts with CD3Z homodimers to complete the TCR-CD3 complex. Alternatively, TCRalpha and TCRbeta can be replaced by TCRgamma and TCRdelta. The CD3E/CD3 epsilon 1-27 peptide also interacts with CD6, NCK1, and NUMB, with the NUMB interaction being crucial for TCR-CD3 internalization and subsequent degradation.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA