

## RGM-C Protein, Cynomolgus (HEK293, His)

Cat. No.:	HY-P701035
Synonyms:	Hemojuvelin; Rgmc; DL-M; HFE2; HFE2AMGC23953; HJV; JH; RGM-C; RGM-C
Species:	Cynomolgus
Source:	HEK293
Accession:	XP_005542141.1 (Q36-D401)
Gene ID:	102138915
Molecular Weight:	50-60 kDa & 30-40 kDa & 20-25 kDa

### PROPERTIES

Biological Activity	Immobilized Cynomolgus RGM-C, His Tag at 0.5µg/ml (100µl/well) on the plate. Dose response curve for Anti-RGM-C Antibody, hFc Tag with the EC <sub>50</sub> of 7.9ng/ml determined by ELISA.
Appearance	Solution.
Formulation	Supplied as a 0.22µm filtered solution of PBS, pH 7.4.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	N/A.
Storage & Stability	Stored at -80°C for 1 year. It is stable at -20°C for 3 months after opening. It is recommended to freeze aliquots at -80°C for extended storage. Avoid repeated freeze-thaw cycles.
Shipping	Shipping with dry ice.

### DESCRIPTION

Background	RGM-C operates as a bone morphogenetic protein (BMP) coreceptor, contributing to the modulation of BMP signaling and playing a crucial role in the regulation of hepcidin (HAMP) expression, thereby influencing iron homeostasis. This protein interacts with BMP2, BMP4, BMP6, BMPR1B, and Tmprss6, forming a network of molecular associations that collectively contribute to its role in mediating BMP-related signaling pathways and iron regulation.
------------	---

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA