

TPBG/5T4 Protein, Human (HEK293, hFc)

Cat. No.:	HY-P701049
Synonyms:	M6P1; 5T4AG; WAIF1; TPBG; 5T4
Species:	Human
Source:	HEK293
Accession:	Q13641 (S32-S355)
Gene ID:	7162
Molecular Weight:	75-105 kDa

PROPERTIES

Biological Activity	Immobilized Human TPBG, hFc Tag at 0.5µg/ml (100µl/Well) on the plate. Dose response curve for Biotinylated Anti-TPBG Antibody, hFc Tag with the EC ₅₀ of 18.5ng/ml determined by ELISA.
Appearance	Solution.
Formulation	Supplied as a 0.22µm filtered solution of PBS, pH 7.4.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	N/A.
Storage & Stability	Stored at -80°C for 1 year. It is stable at -20°C for 3 months after opening. It is recommended to freeze aliquots at -80°C for extended storage. Avoid repeated freeze-thaw cycles.
Shipping	Shipping with dry ice.

DESCRIPTION

Background	The TPBG/5T4 protein appears to serve as an inhibitor of Wnt/beta-catenin signaling, potentially achieved through indirect interaction with LRP6, thereby impeding Wnt3a-dependent LRP6 internalization. This implies a crucial role for TPBG/5T4 in modulating the intricate Wnt/beta-catenin signaling pathway, exerting regulatory influence on cellular responses associated with this pathway. Further exploration of the specific molecular mechanisms governing the interaction between TPBG/5T4 and LRP6, as well as its impact on Wnt3a-dependent LRP6 internalization, could yield valuable insights into the functional significance of TPBG/5T4 in shaping Wnt/beta-catenin signaling dynamics and its potential implications for various cellular processes.
------------	---

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA