

Siglec-10 Protein, Cynomolgus (HEK293, His, solution)

Cat. No.:	HY-P701083
Synonyms:	SLG2; SIGLEC10; MGC126774; PRO940
Species:	Cynomolgus
Source:	HEK293
Accession:	A0A2K5WBX8 (T17-N552)
Gene ID:	102125591
Molecular Weight:	70-80 kDa

PROPERTIES

Biological Activity	Immobilized Cynomolgus Siglec-10, His Tag at 2µg/ml (100µl/well) on the plate. Dose response curve for Anti-Siglec-10 Antibody, hFc Tag with the EC ₅₀ of 42.0ng/ml determined by ELISA.
Appearance	Solution.
Formulation	Supplied as a 0.22µm filtered solution of 25mM MES, 150mM NaCl, 0.5M Arginine, pH 5.0.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	N/A.
Storage & Stability	Stored at -80°C for 1 year. It is stable at -20°C for 3 months after opening. It is recommended to freeze aliquots at -80°C for extended storage. Avoid repeated freeze-thaw cycles.
Shipping	Shipping with dry ice.

DESCRIPTION

Background

Siglec-10 protein, recognized as a putative adhesion molecule, functions in sialic-acid dependent cellular binding, displaying a preference for alpha-2,3- or alpha-2,6-linked sialic acid. The sialic acid recognition site of Siglec-10 may undergo masking due to cis interactions with sialic acids on the same cell surface. In immune responses, it appears to act as an inhibitory receptor, inducing ligand-induced tyrosine phosphorylation and recruiting cytoplasmic phosphatases via their SH2 domains, blocking signal transduction through dephosphorylation of signaling molecules. Siglec-10 is involved in the negative regulation of B-cell antigen receptor signaling, dependent on PTPN6/SHP-1. In association with CD24, it may participate in the selective suppression of the immune response to danger-associated molecular patterns (DAMPs) such as HMGB1, HSP70, and HSP90. Siglec-10, in collaboration with CD24, regulates the immune response of natural killer (NK) cells and plays a role in the control of autoimmunity. During the initiation of adaptive immune responses by CD8-alpha(+) dendritic cells, Siglec-10 inhibits cross-presentation by impairing the formation of MHC class I-peptide complexes, implicating the recruitment of PTPN6/SHP-1 and promoting phagosomal acidification. Siglec-10 interacts with various proteins, including PTPN6/SHP-1, NCF1, CD24, HMGB1, RIGI, CBL, and PTPN11.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA