

CD276/B7-H3 Protein, Cynomolgus (437a.a, HEK293, Fc)

Cat. No.:	HY-P70126
Synonyms:	rCynCD276/B7-H3, Fc; CD276 antigen; CD276; B7 homolog 3; B7-H3; CD276; Cynomolgus CD276 Molecule
Species:	Cynomolgus
Source:	HEK293
Accession:	XP_015308534.1 (L29-E465)
Gene ID:	102131295
Molecular Weight:	100-123 kDa

PROPERTIES

AA Sequence	<pre> LEVQVPEDPV VALVGTDATL RCSFSPEPGF SLAQLNLIWQ LTDTKQLVHS FTEGRDQGSA YANRTALFLD LLAQGNASLR LQVRVVADEG SFTCFVSIIRD FGSAAVSLQV AAPYSKPSMT LEPNKDLRPG DTVTITCSSY RGYPEAEVFW QDGGGAPLTG NVTTSQMANE QGLFDVHSV L RVVLGANGTY SCLVRNPVLQ QDAHGSITIT PQRSP TGAVE VQVPEDPVVA LVGTDATLRC SFSPEPGFSL AQLNLIWQLT DTKQLVHSFT EGRDQGSAYA NRTALFLDLL AQGNASLR LQ RRVVADEGSF TCFVSI RDFG SAAVSLQVAA PYSKPSMTLE PNKDLRPGDT VTITCSSYRG YPEAEVFWQD GQGAPLTGNV TTSQMANEQG LFDVHSVLRV VLGANGTYS C LVRNPVLQQD AHGSVTITGQ PMTFPPE </pre>
Biological Activity	Measured by its binding ability in a functional ELISA. When immobilized Cynomolgus CD276/B7-H3- Fc at 2 µg/mL (100 µl/well), can bind Anti-Human B7-H3 with the ED ₅₀ < 33.31 ng/mL.
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background

The CD276/B7-H3 protein is suggested to play a multifaceted role in the regulation of T-cell-mediated immune responses, potentially acting as a protective factor in tumor cells by inhibiting natural-killer-mediated cell lysis and serving as a marker for the detection of neuroblastoma cells. Additionally, CD276/B7-H3 may be involved in the development of acute and chronic transplant rejection, contributing to the regulation of lymphocytic activity at mucosal surfaces. Notably, it could play a crucial role in providing the placenta and fetus with an immunologically suitable environment throughout pregnancy. Both isoform 1 and isoform 2 of CD276/B7-H3 appear redundant in their ability to modulate CD4 T-cell responses, with isoform 2 demonstrated to enhance the induction of cytotoxic T-cells and selectively stimulate interferon-gamma production in the presence of T-cell receptor signaling. The interaction with TREML2 is identified as enhancing T-cell activation, highlighting the diverse roles CD276/B7-H3 may play in immune regulation and cellular responses.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA