

Screening Libraries

Proteins



Product Data Sheet

FITC-Labeled CD99 Protein, Human (HEK293, Fc)

Cat. No.: HY-P701277

Synonyms: CD99; HBA71; MIC2; MIC2X; MIC2Y; MSK5X; 12E7

Species: Human Source: HEK293

Accession: P14209 (D23-D122)

Gene ID: 4267 Molecular Weight: 45-55 kD

п	п	$\boldsymbol{\smallfrown}$	n	_	п	-		c
-	154		154	-	w	-	ΙE	-
	LIA.3	•		_				_

Appearance	Lyophilized powder.
Formulation	Lyophilized from 0.22 μm filtered solution of PBS, pH7.4. Normally trehalose is added as protectant before lyophilization.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconsititution	It is not recommended to reconstitute to a concentration less than 100 $\mu g/mL$ in ddH ₂ O.
Storage & Stability	Stored at -20°C for 1 year, protect from light. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background

CD99 protein participates in crucial T-cell adhesion processes, contributing to spontaneous rosette formation with erythrocytes. Additionally, it plays a vital role in the late stages of leukocyte extravasation, aiding leukocytes in overcoming the endothelial basement membrane during immune responses. Remarkably, its function is independent of PECAM1, although it acts at the same site. The involvement of CD99 in T-cell adhesion processes underscores its significance in mediating cellular interactions essential for immune function.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA

Page 1 of 1