

FITC-Labeled CD276/B7-H3 Protein, Human (435a.a, HEK293, Fc)

Cat. No.:	HY-P701302
Synonyms:	4lg-B7-H3; B7-H3; CD276; PSEC0249; UNQ309; PRO352; B7 homolog 3
Species:	Human
Source:	HEK293
Accession:	Q5ZPR3 (G27-T461)
Gene ID:	80381
Molecular Weight:	95-115 kDa

PROPERTIES

Biological Activity	Immobilized FITC-Labeled Human B7-H3, hFc Tag at 1 μ g/ml (100 μ l/well) on the plate. Dose response curve for Biotinylated Anti-B7-H3 Antibody, hFc Tag with the EC ₅₀ of 62.5 ng/mL determined by ELISA.
Appearance	Lyophilized powder
Formulation	Lyophilized from 0.22 μ m filtered solution of PBS, pH7.4. Normally trehalose is added as protectant before lyophilization.
Endotoxin Level	<1 EU/ μ g, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 μ g/mL in ddH ₂ O.
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background	<p>The CD276/B7-H3 protein is suggested to play a multifaceted role in the regulation of T-cell-mediated immune responses, potentially acting as a protective factor in tumor cells by inhibiting natural-killer-mediated cell lysis and serving as a marker for the detection of neuroblastoma cells. Additionally, CD276/B7-H3 may be involved in the development of acute and chronic transplant rejection, contributing to the regulation of lymphocytic activity at mucosal surfaces. Notably, it could play a crucial role in providing the placenta and fetus with an immunologically suitable environment throughout pregnancy. Both isoform 1 and isoform 2 of CD276/B7-H3 appear redundant in their ability to modulate CD4 T-cell responses, with isoform 2 demonstrated to enhance the induction of cytotoxic T-cells and selectively stimulate interferon-gamma production in the presence of T-cell receptor signaling. The interaction with TREML2 is identified as enhancing T-cell activation, highlighting the diverse roles CD276/B7-H3 may play in immune regulation and cellular responses.</p>
------------	--

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA