

Neuropilin-1 Protein, Human (619a.a, HEK293, C-His)

| | |
|-------------------|---|
| Cat. No.: | HY-P70147A |
| Synonyms: | rHuNeuropilin-1, His ; Neuropilin-1, CD304; NRP1; NRPNP1; VEGF165R; BDCA4 |
| Species: | Human |
| Source: | HEK293 |
| Accession: | AAH07533.1 (F22-K644) |
| Gene ID: | 8829 |
| Molecular Weight: | approximately 86.92 kDa |

PROPERTIES

AA Sequence

```

FRNDKCGDTI   KIESPGYLT S   PGYPHSYHPS   EKCEWLIQAP
DPYQRIMINF   NPHFDLEDRD   CKYDYVEVFD   GENENGHFRG
KFCGKIAPPP   VVSSGPFLFI   KFVSDYETHG   AGFSIRYEIF
KRGPECSQNY   TTPSGVIKSP   GFPEKYPNSL   ECTYIVFAPK
MSEIILEFES   FDLEPDSNPP   GGMFCRYDRL   EIWDGFPDVG
PHIGRYCGQK   TPGRIRSSSG   ILSMVFYTDS   AIAKEGFSAN
YSVLQSSVSE   DFKCMEALGM   ESGEIHSDQI   TASSQYSTNW
SAERSRLNYP   ENGWTPEGDS   YREWIQVDLG   LLRFVTVAVGT
QGAISKETKK   KYVVKTYKID   VSSNGEDWIT   IKEGNKPVLF
QGNTNPTDVV   VAVFPKPLIT   RFVRIKPATW   ETGISMRFEV
YGCKITDYPC   SGMLGMVSL   ISDSQITSSN   QGDRNWMPEN
IRLVTSRSGW   ALPPAPHSYI   NEWLQIDLGE   EKIVRGI I I Q
GGKHRENKVF   MRKFKIGYSN   NGSDWKIMD   DSKRKAKSFE
GNNNYDTP EL   RTFPALSTRF   IRIYPERATH   GGLGLRME LL
GCEVEAPTAG   PTPPNGNLVD   ECDDDQANCH   SGTGDDFQLT
GGTTVLATEK   PTVIDSTIQS   GIK
  
```

Biological Activity Measured by its ability to inhibit the cell growth of MCF-7 human breast cancer cell line. The ED₅₀ for this effect is 13.44 ng/mL, corresponding to a specific activity is 7.44×10⁴ U/mg.

Appearance Lyophilized powder.

Formulation Lyophilized from a 0.2 μm filtered solution of 20 mM PB, 150 mM NaCl, pH 7.4.

Endotoxin Level <1 EU/μg, determined by LAL method.

Reconstitution It is not recommended to reconstitute to a concentration less than 100 μg/mL in ddH₂O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).

Storage & Stability Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.

Shipping

Room temperature in continental US; may vary elsewhere.

DESCRIPTION**Background**

The Neuropilin-1 protein assumes a significant role as it binds to VEGF-165, potentially inhibiting its binding to cells and inducing apoptosis by sequestering VEGF-165. This dual functionality suggests Neuropilin-1's involvement in the regulation of VEGF-mediated cellular processes. Additionally, Neuropilin-1 may interact with various members of the semaphorin family, indicating its versatility in binding to different ligands. Notably, its expression appears to exert an adverse effect on blood vessel number and integrity, suggesting a potential role in angiogenesis and vascular development. The diverse interactions and regulatory effects of Neuropilin-1 underscore its importance in orchestrating complex cellular processes and its potential implications in angiogenesis and apoptotic pathways.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA