RNF11 Protein, Human

MedChemExpress

| Cat. No.: | HY-P701550 |
|-------------------|-------------------------------|
| Synonyms: | RNF11; RING finger protein 11 |
| Species: | Human |
| Source: | E. coli |
| Accession: | Q9Y3C5 (G2-N154) |
| Gene ID: | 26994 |
| Molecular Weight: | |

| PROPERTIES | |
|-----------------|--|
| Appearance | Solution. |
| Formulation | Supplied as a 0.22 μm filtered solution of 50 mM Tris-HCl, pH7.5, 200 mM NaCl, 20% glycerol. |
| Endotoxin Level | <1 EU/µg, determined by LAL method. |
| Reconsititution | Please use rapid thawing with running water to thaw the protein. |
| | |

| Storage & Stability | Stored at -80°C for 1 year. It is stable at -20°C for 3 months after opening. It is recommended to freeze aliquots at -80°C for | |
|---------------------|---|--|
| | extended storage. Avoid repeated freeze-thaw cycles. | |

| Shipping | Shipping with dry ice. |
|----------|------------------------|
| Simpping | Shipping with dry icc. |

DESCRIPTION

Background RNF11 takes center stage as an essential component of a ubiquitin-editing protein complex, a collaborative ensemble featuring TNFAIP3, ITCH, and TAX1BP1. This complex operates intricately to orchestrate the transient nature of inflammatory signaling pathways, exerting control over TNF- or LPS-mediated activation of NF-kappa-B. RNF11's role extends to promoting the association of TNFAIP3 with RIPK1 post-TNF stimulation. This, in turn, facilitates TNFAIP3's deubiquitination of 'Lys-63' polyubiquitin chains on RIPK1, initiating the formation of 'Lys-48'-polyubiquitin chains that culminate in RIPK1's proteasomal degradation. Furthermore, RNF11 acts as a recruiter, bringing STAMBP to the E3 ubiquitin-ligase SMURF2 for ubiquitination, ultimately resulting in SMURF2's degradation by the 26S proteasome. This intricate web of interactions involves 14-3-3 (when phosphorylated), the E3 ubiquitin-ligases NEDD4, ITCH, SMURF2, and WWP1, as well as the E2 ubiquitin-conjugating enzymes UBE2D1 and UBE2N. Noteworthy associations include those with ZNF350, EPS15, STAMBP, TAX1BP1, TNFAIP3, RIPK1, and GGA1, revealing the multifaceted nature of RNF11's regulatory engagements.

Inhibitors

•

Screening Libraries

•

Proteins

Caution: Product has not been fully validated for medical applications. For research use only.

 Tel: 609-228-6898
 Fax: 609-228-5909
 E-mail: tech@MedChemExpress.com

 Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA