# 

# Product Data Sheet

## MAPKAPK2 Protein, Human

Cat. No.:	HY-P701804
Synonyms:	MAPKAPK2; MAP kinase-activated protein kinase 2; MAPK-activated protein kinase 2; MAPKAP kinase 2; MAPKAP-K2; MAPKAPK-2; MK-2; MK2
Species:	Human
Source:	E. coli
Accession:	P49137-1 (H47-R364, ΔH217-P237, S216G)
Gene ID:	9261
Molecular Weight:	

FROFERILS	
Appearance	Solution.
Formulation	Supplied as a 0.22 $\mu m$ filtered solution of 50 mM Tris-HCl, pH7.5, 200 mM NaCl, 20% glycerol.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconsititution	Please use rapid thawing with running water to thaw the protein.
Storage & Stability	Stored at -80°C for 1 year. It is stable at -20°C for 3 months after opening. It is recommended to freeze aliquots at -80°C for extended storage. Avoid repeated freeze-thaw cycles.
Shipping	Shipping with dry ice.

### DESCRIPTION

# BackgroundMAPKAPK2 protein is a stress-activated serine/threonine-protein kinase with a multifaceted role in cellular processes. It<br/>participates in cytokine production, endocytosis, cytoskeleton reorganization, cell migration, cell cycle control, chromatin<br/>remodeling, DNA damage response, and transcriptional regulation. Upon stress, MAPKAPK2 is phosphorylated and<br/>activated by MAP kinase p38-alpha/MAPK14, leading to the phosphorylation of various substrates. It phosphorylates serine<br/>in specific peptide sequences and targets proteins involved in diverse cellular functions. Notably, MAPKAPK2 mediates the<br/>inflammatory response by regulating the post-transcriptional aspects of tumor necrosis factor (TNF) and IL6 production. It<br/>accomplishes this by phosphorylating AU-rich element (ARE)-binding proteins, affecting mRNA stability and translation.<br/>Additionally, MAPKAPK2 plays a crucial role in the late G2/M checkpoint following DNA damage, contributing to post-<br/>transcriptional mRNA stabilization. Its involvement in the toll-like receptor signaling pathway in dendritic cells underscores<br/>its significance in immune responses. Overall, MAPKAPK2 emerges as a key player in orchestrating cellular responses to<br/>stress and various signaling pathways.

### Caution: Product has not been fully validated for medical applications. For research use only.

 Tel: 609-228-6898
 Fax: 609-228-5909
 E-mail: tech@MedChemExpress.com

 Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA