Proteins



Product Data Sheet

GPR87 Protein, Human (Cell-Free, Strep)

Cat. No.: HY-P702309

Synonyms: G-protein coupled receptor 87; G-protein coupled receptor 95

Species:

E. coli Cell-free Source: Q9BY21 (M1-V358) Accession:

Gene ID: 53836 44.5 kDa Molecular Weight:

PROPERTIES

	_		
$\Lambda \Lambda$	Sea	HAN	00

MGFNLTLAKL PNNELHGQES HNSGNRSDGP GKNTTLHNEF DTIVLPVLYL IIFVASILLN GLAVWIFFHI RNKTSFIFYL KNIVVADLIM TLTFPFRIVH DAGFGPWYFK FILCRYTSVL FYANMYTSIV FLGLISIDRY LKVVKPFGDS RMYSITFTKV LSVCVWVIMA VLSLPNIILT NGQPTEDNIH DCSKLKSPLG VKWHTAVTYV $\mathsf{N}\;\mathsf{S}\;\mathsf{C}\;\mathsf{L}\;\mathsf{F}\;\mathsf{V}\;\mathsf{A}\;\mathsf{V}\;\mathsf{L}\;\mathsf{V}$ ILIGCYIAIS RYIHKSSRQF ISQSSRKRKH NQSIRVVAV FFTCFLPYHL CRIPFTFSHL ILYYCKEITL DRLLDESAQK FLSACNVCLD PIIYFFMCRS FSRRLFKKSN IRTRSESIRS LQSVRRSEVR IYYDYTDV

Appearance

Lyophilized powder.

Formulation

Lyophilized from a 0.22 µm filtered solution of Tris/PBS-based buffer, 6% Trehalose, pH 8.0.

Endotoxin Level

<1 EU/µg, determined by LAL method.

Reconsititution

It is not recommended to reconstitute to a concentration less than $100 \, \mu g/mL$ in ddH_2O . For long term storage it is recommended to add 5-50% of glycerol (final concentration). Our default final concentration of glycerol is 50%. Customers could use it as reference.

Storage & Stability

Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.

Shipping

Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background

GPR87, serving as a receptor for lysophosphatidic acid (LPA), plays a crucial role in p53/TP53-dependent survival pathways activated in response to DNA damage. The receptor's engagement with LPA is instrumental in mediating cellular responses

		medical applications. F	
Tel: 609-228-6	Fax: 609-228-5909 Park Dr, Suite Q, Monr		edChemExpress.com 52, USA
		E-mail: tech@M mouth Junction, NJ 088	

to DNA damage, emphasizing its significance in processes related to cell survival and DNA damage repair.

Page 2 of 2 www.MedChemExpress.com