

GST Protein, Schistosoma Japonicum

Cat. No.:	HY-P70270
Synonyms:	rSchGlutathione S-transferase class-mu 26 kDa isozyme; Glutathione S-transferase class-mu 26 kDa isozyme; GST 26; Sj26 antigen; SjGST
Species:	Others
Source:	E. coli
Accession:	P08515 (M1-K218)
Gene ID:	/
Molecular Weight:	Approximately 28.0 kDa

PROPERTIES

AA Sequence	<pre> M S P I L G Y W K I K G L V Q P T R L L L E Y L E E K Y E E H L Y E R D E G D K W R N K K F E L G L E F P N L P Y Y I D G D V K L T Q S M A I I R Y I A D K H N M L G G C P K E R A E I S M L E G A V L D I R Y G V S R I A Y S K D F E T L K V D F L S K L P E M L K M F E D R L C H K T Y L N G D H V T H P D F M L Y D A L D V V L Y M D P M C L D A F P K L V C F K K R I E A I P Q I D K Y L K S S K Y I A W P L Q G W Q A T F G G G D H P P K </pre>
Biological Activity	Data is not available.
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O.
Storage & Stability	Stored at -20°C. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer. It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA