

## Product Data Sheet

## Grancalcin/GCA Protein, Human (His)

Cat. No.:	HY-P70334
Synonyms:	rHuGrancalcin/GCA; Grancalcin; GCA; GCL
Species:	Human
Source:	E. coli
Accession:	P28676 (M1-I217)
Gene ID:	25801
Molecular Weight:	Approximately 25 kDa

PROPERTIES		
PROPERTIES		
AA Sequence	MAYPGYGGGFGNFSIQVPGMQMGQPVPETGPAILLDGYSGPAYSDTYSSAGDSVYTYFSAVAGQDGEVDAEELQRCLTQSGINGTYSPFSLETCRIMIAMLDRDHTGKMGFNAFKELWAALNAWKENFMTVDQDGSGTVEHHELRQAIGLMGYRLSPQTLTTIVKRYSKNGRIFFDDYVACCVKLRALTDFFRKRDHLQQGSANFIYDDFLQGTMAI	
Biological Activity	Data is not available.	
Appearance	Lyophilized powder.	
Formulation	Lyophilized from a 0.2 $\mu m$ filtered solution of 50 mM Tris-HCL, 200 mM NaCl, 100 mM arginine, pH 8.0.	
Endotoxin Level	<1 EU/µg, determined by LAL method.	
Reconsititution	It is not recommended to reconstitute to a concentration less than 100 μg/mL in ddH <sub>2</sub> O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).	
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.	
Shipping	Room temperature in continental US;may vary elsewhere.	

## **DESCRIPTION Background** Grancalcin (GCA) is a calcium-binding protein with putative functions in neutrophil adhesion to fibronectin and the formation of focal adhesions. Its role in calcium binding suggests involvement in signaling pathways that regulate cellular adhesion processes. GCA is known to form homodimers and has interactions with SRI and LCP1, further implicating its participation in protein-protein interactions that could modulate cellular functions. It has to emphasize GCA's potential

## Caution: Product has not been fully validated for medical applications. For research use only.

 Tel: 609-228-6898
 Fax: 609-228-5909
 E-mail: tech@MedChemExpress.com

 Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA