

## CD28 Protein, Human/Cynomolgus (HEK293, His)

Cat. No.:	HY-P70486
Synonyms:	CD28; CD28 antigen; CD28 molecule; T-cell-specific surface glycoprotein CD28; Tp44; TP44
Species:	Human
Source:	HEK293
Accession:	P10747 (N19-P152)
Gene ID:	940
Molecular Weight:	30-40 kDa

### PROPERTIES

AA Sequence	<div> <div>N K I L V K Q S P M</div> <div>L D S A V E V C V V</div> <div>L Q N L Y V N Q T D</div> <div>H L C P S P L F P G</div> </div> <div> <div>L V A Y D N A V N L</div> <div>Y G N Y S Q Q L Q V</div> <div>I Y F C K I E V M Y</div> <div>P S K P</div> </div> <div> <div>S C K Y S Y N L F S</div> <div>Y S K T G F N C D G</div> <div>P P P Y L D N E K S</div> </div> <div> <div>R E F R A S L H K G</div> <div>K L G N E S V T F Y</div> <div>N G T I I H V K G K</div> </div>
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 µm filtered solution of 20 mM Tris, 150 mM NaCl, pH 8.0.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH <sub>2</sub> O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

### DESCRIPTION

Background	<p>CD28 protein plays a pivotal role in T-cell activation, promoting cell proliferation, cytokine production, and T-cell survival. Upon ligation with TCR/CD3 and CD40L costimulation, CD28 enhances the production of IL4 and IL10 in T-cells, contributing to immune response modulation. Additionally, isoform 3 of CD28 facilitates CD40L-mediated activation of NF-kappa-B and kinases MAPK8 and PAK2 in T-cells. The protein forms homodimers through disulfide linkages and interacts with various molecules, including DUSP14, CD80/B7-1, CD86/B7-2/B70, and GRB2. Isoform 3 specifically interacts with CD40LG, highlighting its multifaceted role in mediating immune responses and cellular signaling pathways.</p>
------------	--

---

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA