

VEGF-C Protein, Human (196a.a, HEK 293, His)

Cat. No.:	HY-P70501
Synonyms:	Vascular Endothelial Growth Factor C; VEGF-C; Flt4 Ligand; Flt4-L; Vascular Endothelial Growth Factor-Related Protein; VRP; VEGFC
Species:	Human
Source:	HEK 293
Accession:	P49767 (F32-R227)
Gene ID:	7424
Molecular Weight:	25-30 kDa

PROPERTIES

AA Sequence	<pre> F E S G L D L S D A E P D A G E A T A Y A S K D L E E Q L R S V S S V D E L M T V L Y P E Y W K M Y K C Q L R K G G W Q H N R E Q A N L N S R T E E T I K F A A A H Y N T E I L K S I D N E W R K T Q C M P R E V C I D V G K E F G V A T N T F F K P P C V S V Y R C G G C C N S E G L Q C M N T S T S Y L S K T L F E I T V P L S Q G P K P V T I S F A N H T S C R C M S K L D V Y R Q V H S I I R R </pre>
Biological Activity	Data is not available.
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 µm filtered solution of 20 mM PB, 150 mM NaCl, pH 8.0.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O.
Storage & Stability	Stored at -20°C. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer. It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA