

## FLT3LG Protein, Human (HEK 293, His)

Cat. No.:	HY-P70544
Synonyms:	Fms-Related Tyrosine Kinase 3 Ligand; Flt3 Ligand; Flt3L; SL Cytokine; FLT3LG
Species:	Human
Source:	HEK 293
Accession:	P49771 (T27-P184)
Gene ID:	2323
Molecular Weight:	24-32 kDa

### PROPERTIES

AA Sequence	<p>T Q D C S F Q H S P    I S S D F A V K I R    E L S D Y L L Q D Y    P V T V A S N L Q D</p> <p>E E L C G G L W R L    V L A Q R W M E R L    K T V A G S K M Q G    L L E R V N T E I H</p> <p>F V T K C A F Q P P    P S C L R F V Q T N    I S R L L Q E T S E    Q L V A L K P W I T</p> <p>R Q N F S R C L E L    Q C Q P D S S T L P    P P W S P R P L E A    T A P T A P Q P</p>
Biological Activity	Data is not available.
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 µm filtered solution of 20 mM Tris-HCl, 150 mM NaCl, pH 8.0.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH <sub>2</sub> O.
Storage & Stability	Stored at -20°C. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer. It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA