

## TNF-alpha/TNFSF2 Protein, Mouse (147a.a)

<b>Cat. No.:</b>	HY-P70571
<b>Synonyms:</b>	Tumor Necrosis Factor; Cachectin; TNF-Alpha; Tumor Necrosis Factor Ligand Superfamily Member 2; TNF-a; Tumor Necrosis Factor; Membrane Form; Tumor Necrosis Factor; Soluble Form; Tnf; Tnfa; Tnfsf2
<b>Species:</b>	Mouse
<b>Source:</b>	E. coli
<b>Accession:</b>	P06804 (D89-L235)
<b>Gene ID:</b>	21926
<b>Molecular Weight:</b>	Approximately 17.0 kDa

### PROPERTIES

<b>AA Sequence</b>	<p>           D K P V A H V V A N    H Q V E E Q L E W L    S Q R A N A L L A N    G M D L K D N Q L V            V P A D G L Y L V Y    S Q V L F K G Q G C    P D Y V L L T H T V    S R F A I S Y Q E K            V N L L S A V K S P    C P K D T P E G A E    L K P W Y E P I Y L    G G V F Q L E K G D            Q L S A E V N L P K    Y L D F A E S G Q V    Y F G V I A L         </p>
<b>Biological Activity</b>	The cytotoxicity assay using L-929 mouse fibroblast cells in the presence of the metabolic inhibitor actinomycin D has an ED <sub>50</sub> value of 2-8 pg/mL.
<b>Appearance</b>	Lyophilized powder
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4 or 50 mM Tris-HCL, 300 mM NaCl, pH 7.4.
<b>Endotoxin Level</b>	<1 EU/µg, determined by LAL method.
<b>Reconstitution</b>	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH <sub>2</sub> O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).
<b>Storage &amp; Stability</b>	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
<b>Shipping</b>	Room temperature in continental US; may vary elsewhere.

### DESCRIPTION

<b>Background</b>	<p>TNF alpha is produced by various types of cells including macrophages, monocytes, neutrophils, T cells, and NK-cells<sup>[2]</sup>. The amino acid sequence of human TNF alpha protein has low homology between mouse, rat, bovine, cynomolgus TNF alpha protein. While, human TNF alpha shares 94.85% aa sequence identity with cynomolgus TNF alpha protein, mouse TNF alpha shares 94.47% aa sequence identity with rat TNF alpha protein.</p> <p>TNF alpha exists in two forms; a type II transmembrane protein (tmTNF-α) and a mature soluble protein (sTNF-α). TNF-α</p>
-------------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

---

binds to its receptors, mainly TNFR1 and TNFR2, and then transmits molecular signals for biological functions such as inflammation and cell death. Both sTNF- $\alpha$  and tmTNF- $\alpha$  activate TNFR1, and process a death domain (DD) that interacts with the TNFR1-associated death domain (TRADD) adaptor protein. The TNFR2 signaling pathway is mainly activated by tmTNF- $\alpha$ . TNFR1 signaling tends to be pro-inflammatory and apoptotic. TNFR2 results in NF- $\kappa$ B and MAPKs and AKT activation, TNFR2 activation is associated with homeostatic bioactivities such as tissue regeneration, cell proliferation, and cell survival, as well as host defense and inflammation<sup>[1]</sup>.

TNF-alpha is critical for normal immune response, abnormal secretion TNF alpha activates synovial fibroblasts, keratinocytes, osteoclasts, induces rheumatoid arthritis, inflammatory bowel disease, psoriatic arthritis (PsA), and noninfectious uveitis (NIU)<sup>[3]</sup>. TNF alpha positively regulates endogenous TNF- $\alpha$  expression levels independently of Pgp efflux activity, induces IHF cells proliferation<sup>[4]</sup>. TNF alpha in tissues may promote cancer growth, invasion, and metastasis. Besides, TNF alpha stimulates NF- $\kappa$ B pathway via TNFR2 and anti-TNF- $\alpha$  MAb significantly suppresses the tumor development in colitis-associated cancer (CAC) mouse<sup>[5]</sup>. TNF alpha as a proneurogenic factor activates the SAPK/JNK pathway and can facilitate neuronal replacement and brain repair in response to brain injury<sup>[6]</sup>.

---

## REFERENCES

- [1]. Horiuchi T, et al. Transmembrane TNF-alpha: structure, function and interaction with anti-TNF agents. *Rheumatology (Oxford)*. 2010 Jul;49(7):1215-28.
- [2]. El-Tahan RR, et al. TNF- $\alpha$  gene polymorphisms and expression. *Springerplus*. 2016 Sep 7;5(1):1508.
- [3]. Jang DI, et al. The Role of Tumor Necrosis Factor Alpha (TNF- $\alpha$ ) in Autoimmune Disease and Current TNF- $\alpha$  Inhibitors in Therapeutics. *Int J Mol Sci*. 2021 Mar 8;22(5):2719.
- [4]. Berguetti T, et al. TNF- $\alpha$  Modulates P-Glycoprotein Expression and Contributes to Cellular Proliferation via Extracellular Vesicles. *Cells*. 2019 May 24;8(5):500.
- [5]. Onizawa M, et al. Signaling pathway via TNF-alpha/NF-kappaB in intestinal epithelial cells may be directly involved in colitis-associated carcinogenesis. *Am J Physiol Gastrointest Liver Physiol*. 2009 Apr;296(4):G850-9.
- [6]. Bernardino L, et al. Tumor necrosis factor-alpha modulates survival, proliferation, and neuronal differentiation in neonatal subventricular zone cell cultures. *Stem Cells*. 2008 Sep;26(9):2361-71.
- [7]. Matsuno H, et al. The role of TNF-alpha in the pathogenesis of inflammation and joint destruction in rheumatoid arthritis (RA): a study using a human RA/SCID mouse chimera. *Rheumatology (Oxford)*. 2002 Mar;41(3):329-37.

---

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA