

IFN-gamma Protein, Human (HEK293)

Cat. No.:	HY-P70610
Synonyms:	Interferon Gamma; IFN-Gamma; Immune Interferon; IFNG
Species:	Human
Source:	HEK293
Accession:	P01579 (Q24-Q166)
Gene ID:	3458
Molecular Weight:	20-25 & 16-17kDa

PROPERTIES

AA Sequence	<p>Q D P Y V K E A E N L K K Y F N A G H S D V A D N G T L F L G I L K N W K E E S</p> <p>D R K I M Q S Q I V S F Y F K L F K N F K D D Q S I Q K S V E T I K E D M N V K</p> <p>F F N S N K K K R D D F E K L T N Y S V T D L N V Q R K A I H E L I Q V M A E L</p> <p>S P A A K T G K R K R S Q M L F R G R R A S Q</p>
Biological Activity	Measured by its ability to inhibit the proliferation of HT-29 human coloncancer cells. The ED ₅₀ for this effect is <0.3176 ng/mL, corresponding to a specific activity is >3.148×10 ⁶ Unit/mg.
Appearance	Lyophilized powder
Formulation	Lyophilized from a 0.2 μm filtered solution of 20 mM PB, 4% Mannitol, 2% Sucrose, 0.02% Tween80, pH 7.4 or PBS, pH 7.4.
Endotoxin Level	<1 EU/μg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 μg/mL in ddH ₂ O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background	IFN-gamma is a dimeric soluble cytokine that is the only member of type II interferon IFN-gamma is produced by immune cells T cells and NK cells and plays an important role in antimicrobial, antiviral and anti-tumor responses by activating effector immune cells and enhancing antigen presentation. IFN-gamma influences gene regulation by interacting with its receptor IFNGR1 through the JAK-STAT pathway, and can also trigger mTOR, MAPK, and PI3K/AKT signaling pathways. IFN-gamma plays a role in the Class I antigen presentation pathway by inducing the substitution of the catalytic proteasome
------------	--

subunit for the immune proteasome subunit. IFN-gamma upregulates the MHC II complex on the cell surface by promoting the expression of several key molecules such as pepsin B/CTSB, H/CTSH, and L/CTSL. IFN-gamma is involved in the regulation of hematopoietic stem cells under developmental and homeostasis conditions by influencing the development, quiescence and differentiation of hematopoietic stem cells^{[1][2][3][4][5]}.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA