

IL-2R beta/CD122 Protein, Human (HEK 293, His)

Cat. No.:	HY-P70631
Synonyms:	IL-2R beta; IL-2 R beta; IL-2RB; Interleukin-2 receptor subunit beta; IL2RB; IL-2 receptor subunit beta; IL-2R subunit beta; High affinity IL-2 receptor subunit beta; CD122
Species:	Human
Source:	HEK 293
Accession:	P14784 (A27-D239)
Gene ID:	3560
Molecular Weight:	35-40 kDa

PROPERTIES

AA Sequence	<pre> A V N G T S Q F T C F Y N S R A N I S C V W S Q D G A L Q D T S C Q V H A W P D R R R W N Q T C E L L P V S Q A S W A C N L I L G A P D S Q K L T T V D I V T L R V L C R E G V R W R V M A I Q D F K P F E N L R L M A P I S L Q V V H V E T H R C N I S W E I S Q A S H Y F E R H L E F E A R T L S P G H T W E E A P L L T L K Q K Q E W I C L E T L T P D T Q Y E F Q V R V K P L Q G E F T T W S P W S Q P L A F R T K P A A L G K D </pre>
Biological Activity	Data is not available.
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O.
Storage & Stability	Stored at -20°C. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer. It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA