

VEGFR-1 Protein, Human (730a.a, HEK293, Fc)

Cat. No.:	HY-P70636
Synonyms:	Vascular endothelial growth factor receptor 1; VEGFR-1; Fms-like tyrosine kinase 1; FLT-1; Tyrosine-protein kinase FRT; Tyrosine-protein kinase receptor FLT; Vascular permeability factor receptor
Species:	Human
Source:	HEK293
Accession:	P17948 (S27-N756)
Gene ID:	2321
Molecular Weight:	150-175 kDa

PROPERTIES

AA Sequence

S	K	L	K	D	P	E	L	S	L	K	G	T	Q	H	I	M	Q	A	G	Q	T	L	H	L	Q	C	R	G	E	A	A	H	K	W	S	L	P	E	M
V	S	K	E	S	E	R	L	S	I	T	K	S	A	C	G	R	N	G	K	Q	F	C	S	T	L	T	L	N	T	A	Q	A	N	H	T	G	F	Y	S
C	K	Y	L	A	V	P	T	S	K	K	K	E	T	E	S	A	I	Y	I	F	I	S	D	T	G	R	P	F	V	E	M	Y	S	E	I	P	E	I	
H	M	T	E	G	R	E	L	V	I	P	C	R	V	T	S	P	N	I	T	V	T	L	K	K	F	P	L	D	T	L	I	P	D	G	K	R	I	I	W
D	S	R	K	G	F	I	I	S	N	A	T	Y	K	E	I	G	L	L	T	C	E	A	T	V	N	G	H	L	Y	K	T	N	Y	L	T	H	R	Q	T
N	T	I	I	D	V	Q	I	S	T	P	R	P	V	K	L	L	R	G	H	T	L	V	L	N	C	T	A	T	T	P	L	N	T	R	V	Q	M	T	W
S	Y	P	D	E	K	N	K	R	A	S	V	R	R	R	I	D	Q	S	N	S	H	A	N	I	F	Y	S	V	L	T	I	D	K	M	Q	N	K	D	K
G	L	Y	T	C	R	V	R	S	G	P	S	F	K	S	V	N	T	S	V	H	I	Y	D	K	A	F	I	T	V	K	H	R	K	Q	Q	V	L	E	T
V	A	G	K	R	S	Y	R	L	S	M	K	V	K	A	F	P	S	P	E	V	V	W	L	K	D	G	L	P	A	T	E	K	S	A	R	Y	L	T	R
G	Y	S	L	I	I	K	D	V	T	E	E	D	A	G	N	Y	T	I	L	L	S	I	K	Q	S	N	V	F	K	N	L	T	A	T	L	I	V	N	V
K	P	Q	I	Y	E	K	A	V	S	S	F	P	D	P	A	L	Y	P	L	G	S	R	Q	I	L	T	C	T	A	Y	G	I	P	Q	P	T	I	K	W
F	W	H	P	C	N	H	N	H	S	E	A	R	C	D	F	C	S	N	N	E	E	S	F	I	L	D	A	D	S	N	M	G	N	R	I	E	S	I	T
Q	R	M	A	I	I	E	G	K	N	K	M	A	S	T	L	V	V	A	D	S	R	I	S	G	I	Y	I	C	I	A	S	N	K	V	G	T	V	G	R
N	I	S	F	Y	I	T	D	V	P	N	G	F	H	V	N	L	E	K	M	P	T	E	G	E	D	L	K	L	S	C	T	V	N	K	F	L	Y	R	D
V	T	W	I	L	L	R	T	V	N	N	R	T	M	H	Y	S	I	S	K	Q	K	M	A	I	T	K	E	H	S	I	T	L	N	L	T	I	M	N	V
S	L	Q	D	S	G	T	Y	A	C	R	A	R	N	V	Y	T	G	E	E	I	L	Q	K	K	E	I	T	I	R	D	Q	E	A	P	Y	L	L	R	N
L	S	D	H	T	V	A	I	S	S	S	T	T	L	D	C	H	A	N	G	V	P	E	P	Q	I	T	W	F	K	N	N	H	K	I	Q	Q	E	P	G
I	I	L	G	P	G	S	S	T	L	F	I	E	R	V	T	E	E	D	E	G	V	Y	H	C	K	A	T	N	Q	K	G	S	V	E	S	S	A	Y	L
T	V	Q	G	T	S	D	K	S	N																														

Biological Activity The enzyme activity of this recombinant protein is testing in progress, we cannot offer a guarantee yet.

Appearance Lyophilized powder.

Formulation Lyophilized from a 0.2 µm filtered solution of 20 mM PB, 150 mM NaCl, 5% Trehalose, 5% Mannitol, 0.01% Tween 80, pH 7.0.

Endotoxin Level <1 EU/µg, determined by LAL method.

Reconstitution It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH₂O.

Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background

VEGFR-1 Protein is a tyrosine-protein kinase that serves as a cell-surface receptor for VEGFA, VEGFB, and PGF. It plays a crucial role in various biological processes, including embryonic vasculature development, angiogenesis regulation, cell survival, cell migration, macrophage function, chemotaxis, and cancer cell invasion. Additionally, VEGFR-1 acts as a positive regulator of postnatal retinal hyaloid vessel regression and may function as a negative regulator of embryonic angiogenesis by inhibiting excessive endothelial cell proliferation. In adulthood, it promotes endothelial cell proliferation, survival, and angiogenesis, although its proliferative effects appear to be cell-type specific. VEGFR-1 has a high affinity for VEGFA and acts as a negative regulator of its signaling by limiting the availability of free VEGFA and preventing its binding to KDR. It modulates KDR signaling by forming heterodimers with KDR. Ligand binding to VEGFR-1 activates several signaling cascades, including PLCG, MAPK1/ERK2, MAPK3/ERK1, AKT1, and PI3K pathways. VEGFR-1 also phosphorylates SRC, YES1, CBL, AKT1, PTK2/FAK1, and PLCG.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA