

IL-2 Protein, Mouse (His)

Cat. No.:	HY-P70646
Synonyms:	aldesleukin; interleukin 2; interleukin-2; IL-2; IL2; T-cell growth factor; T cell growth factor; TCGF
Species:	Mouse
Source:	E. coli
Accession:	P04351
Gene ID:	16183
Molecular Weight:	Approximately 17.0 kDa

PROPERTIES

AA Sequence	<p>A P T S S S T S S S T A E A Q Q Q Q Q Q Q Q Q Q Q Q H L E Q L L M D L Q E L L S</p> <p>R M E N Y R N L K L P R M L T F K F Y L P K Q A T E L K D L Q C L E D E L G P L</p> <p>R H V L D L T Q S K S F Q L E D A E N F I S N I R V T V V K L K G S D N T F E C</p> <p>Q F D D E S A T V V D F L R R W I A F C Q S I I S T S P Q</p>
Biological Activity	The cell proliferation assay using CTLL-2 mouse cytotoxic T cells has a specific activity of $\geq 1 \times 10^7$ IU/mg.
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 μ m filtered solution of 20 mM Sodium Citrate, 0.2% Tween 80, pH 3.0.
Endotoxin Level	<1 EU/ μ g, determined by LAL method.
Reconstitution	Reconstitute at 100 μ g/mL in ddH ₂ O.
Storage & Stability	Stored at -20°C. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer. It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

REFERENCES

[1]. Abul K Abbas, et al. Revisiting IL-2: Biology and therapeutic prospects. Sci Immunol. 2018 Jul 6;3(25):eaat1482.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA