

NKG2D/CD314 Protein, Human (HEK293, His)

Cat. No.:	HY-P70688
Synonyms:	CD314; KLRK1; CD314 antigen; Killer cell lectin-like receptor subfamily K member 1; killer cell lectin-like receptor subfamily K; member 1; KLR; NK cell receptor D; NKG2-D; NKG2-D type II integral membrane protein; NKG2-D-activating NK receptor; NKG2D
Species:	Human
Source:	HEK293
Accession:	P26718 (F78-V216)
Gene ID:	22914
Molecular Weight:	20-30 kDa

PROPERTIES

AA Sequence	<p> F L N S L F N Q E V Q I P L T E S Y C G P C P K N W I C Y K N N C Y Q F F D E S K N W Y E S Q A S C M S Q N A S L L K V Y S K E D Q D L L K L V K S Y H W M G L V H I P T N G S W Q W E D G S I L S P N L L T I I E M Q K G D C A L Y A S S F K G Y I E N C S T P N T Y I C M Q R T V </p>
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background	<p>NKG2D/CD314 protein operates as an activating and costimulatory receptor essential for immunosurveillance, binding to diverse cellular stress-inducible ligands presented on autologous tumor cells and virus-infected cells. It plays a dual role in innate immune responses, stimulating both activating killer (NK) cells and acting as a costimulatory receptor for T-cell receptors (TCR) in CD8(+) T-cell-mediated adaptive immune responses, enhancing T-cell activation. The receptor facilitates perforin-mediated elimination of ligand-expressing tumor cells, and its signaling cascades involve calcium influx, ultimately leading to TNF-alpha expression. Additionally, NKG2D/CD314 participates in NK cell-mediated bone marrow graft rejection and may regulate the differentiation and survival of NK cells. Its ligand-binding capacity extends to various subfamilies of MHC class I-related glycoproteins, including MICA, MICB, RAET1E, RAET1G, RAET1L/ULBP6, ULBP1, ULBP2, ULBP3</p>
-------------------	--

(ULBP2>ULBP1>ULBP3), and ULBP4. The protein forms homodimers through disulfide linkage and heterohexamers with HCST/DAP10 subunits, a crucial interaction for NK cell surface expression and cytotoxicity induction. Furthermore, it can establish disulfide-bonded heterodimers with CD94 and interacts with CEACAM1, recruiting PTPN6 for VAV1 dephosphorylation, while not interacting with TYROBP.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA