

Certificate of Analysis

Fc gamma RIIIA/CD16a Protein, Human (HEK293, His)

| Cat. No.: | HY-P70705 |
|-------------------|-------------------|
| Batch No.: | 272763 |
| Species: | Human |
| Source: | HEK293 |
| Tag: | C-6*His |
| Accession: | P08637 (G17-Q208) |
| Gene ID: | 2214 |
| Molecular Weight: | 36-44 kDa |

ANALYTICAL DATA

| TEST | Specifications | Results |
|---------------------|--|-------------|
| Purity | Greater than 95% as determined by reducing SDS-PAGE. | ≥95.0% |
| Endotoxin Level | <1 EU/µg, determined by LAL method. | <1 EU/µg |
| Biological Activity | Measured by its binding ability in a functional ELISA. Immobilized Human Fc γ RIIIA / CD16a recombinant protein at 2 μ g/mL(100 μ L/well) can bind Human IgG1. The ED ₅₀ for this effect is 490.6 ng/mL, corresponding to a specific activity is 2238.3 Unit/mg. | 490.6 ng/mL |
| Appearance | Lyophilized powder. | |
| Reconstitution | It is not recommended to reconstitute to a concentration less than 100 μg/mL in ddH ₂ O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose). | |
| Storage & Stability | Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or - 20°C for longer. It is recommended to freeze aliquots at -20°C or -80°C for extended storage. | |
| Shipping | Room temperature in continental US; may vary elsewhere. | |

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

28-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA