

## IL-5 Protein, Mouse (HEK293, His)

Cat. No.:	HY-P70746
Synonyms:	Interleukin-5; IL-5; B-cell differentiation factor I; Eosinophil differentiation factor; T-cell replacing factor; TRF; IL5
Species:	Mouse
Source:	HEK293
Accession:	P04401 (M21-G133)
Gene ID:	16191
Molecular Weight:	14-26 kDa

### PROPERTIES

AA Sequence	ME I P M S T V V K      E T L T Q L S A H R      A L L T S N E T M R      L P V P T H K N H Q L C I G E I F Q G L      D I L K N Q T V R G      G T V E M L F Q N L      S L I K K Y I D R Q K E K C G E E R R R      T R Q F L D Y L Q E      F L G V M S T E W A      M E G
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH <sub>2</sub> O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

### DESCRIPTION

Background	IL-5 Protein, a homodimeric cytokine predominantly expressed by T-lymphocytes and NK cells, plays a pivotal role in the regulation of eosinophils by influencing their survival, differentiation, and chemotaxis. Additionally, IL-5 acts on both activated and resting B-cells, stimulating immunoglobulin production, growth, and differentiation. Mechanistically, the biological effects of IL-5 are mediated through a receptor complex composed of the IL5RA subunit and the cytokine receptor common subunit beta/CSF2RB. Upon binding to the receptor, IL-5 triggers the activation of various kinases, including LYN, SYK, and JAK2, thus propagating signals through the RAS-MAPK and JAK-STAT5 pathways (By similarity). Structurally, IL-5 forms a homodimer that is disulfide-linked and interacts with its receptor components IL5RA and CSF2RB, highlighting the specificity and complexity of its molecular interactions in orchestrating immune responses.
------------	---

---

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA