

## IL-2 Protein, Human (HEK293, His)

Cat. No.:	HY-P70758
Synonyms:	Interleukin-2; IL-2; T-Cell Growth Factor; TCGF; Aldesleukin; IL2
Species:	Human
Source:	HEK293
Accession:	P60568 (A21-T153)
Gene ID:	3558
Molecular Weight:	14-18 kDa

### PROPERTIES

AA Sequence	<pre> A P T S S S T K K T   Q L Q L E H L L L D   L Q M I L N G I N N   Y K N P K L T R M L T F K F Y M P K K A   T E L K H L Q C L E   E E L K P L E E V L   N L A Q S K N F H L R P R D L I S N I N   V I V L E L K G S E   T T F M C E Y A D E   T A T I V E F L N R W I T F C Q S I I S   T L T           </pre>
Biological Activity	The cell proliferation assay using CTLL-2 mouse cytotoxic T cells has a specific activity of $\geq 1 \times 10^7$ IU/mg.
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 $\mu$ m filtered solution of 20 mM PB, 150 mM NaCl, pH 7.4.
Endotoxin Level	<1 EU/ $\mu$ g, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 $\mu$ g/mL in ddH <sub>2</sub> O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

### DESCRIPTION

Background	<p>The cytokine interleukin-2 (IL-2), produced primarily by activated CD4-positive helper T-cells and, to a lesser extent, by activated CD8-positive T-cells and natural killer (NK) cells, plays pivotal roles in the immune response and tolerance. IL-2 binds to a receptor complex composed of either the high-affinity trimeric IL-2R (IL2RA/CD25, IL2RB/CD122, and IL2RG/CD132) or the low-affinity dimeric IL-2R (IL2RB and IL2RG). This interaction induces oligomerization and conformational changes in the IL-2R subunits, initiating downstream signaling with the phosphorylation of JAK1 and JAK3. Subsequently, JAK1 and JAK3 phosphorylate the receptor, creating a docking site for the phosphorylation of various</p>
------------	---

---

substrates, including STAT5. This process activates multiple pathways, including STAT, phosphoinositide-3-kinase/PI3K, and mitogen-activated protein kinase/MAPK pathways. IL-2 functions as a T-cell growth factor, enhances NK-cell cytolytic activity, and promotes robust proliferation of activated B-cells, leading to increased immunoglobulin production. Furthermore, IL-2 plays a crucial role in regulating the adaptive immune system by controlling the survival and proliferation of regulatory T-cells, essential for maintaining immune tolerance. Additionally, IL-2 participates in the differentiation and homeostasis of various effector T-cell subsets, including Th1, Th2, Th17, as well as memory CD8-positive T-cells.

---

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA