

IgG1 Protein, Human (D239E, L241M, HEK293)

Cat. No.:	HY-P70796
Synonyms:	Ig gamma-1 chain C region; IGHG1
Species:	Human
Source:	HEK293
Accession:	P01857-1 (D104-K330, D239E, L241M)
Gene ID:	3500
Molecular Weight:	Approximately 33 kDa

PROPERTIES

AA Sequence	<pre> DKTHTCPPCP APELLGGPSV FLFPPKPKDT LMISRTPEVT CVVVDVSHED PEVKFNWYVD GVEVHNAKTK PREEQYNSTY RVVSVLTVHL QDWLNGKEYK CKVSNKALPA PIEKTISKAK GQPREPQVYT LPPSREEMTK NQVSLTCLVK GFYPSDIAVE WESNGQPENN YKTTTPVLDL DGSFFLYSKL TVDKSRWQQG NVFSCSVMHE ALHNNHYTKS LSLSPGK </pre>
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background	<p>The constant region of immunoglobulin heavy chains, known as antibodies, represents membrane-bound or secreted glycoproteins produced by B lymphocytes. In the recognition phase of humoral immunity, these membrane-bound immunoglobulins act as receptors, initiating the clonal expansion and differentiation of B lymphocytes into immunoglobulin-secreting plasma cells upon binding specific antigens. Secreted immunoglobulins play a crucial role in the effector phase of humoral immunity, leading to the elimination of bound antigens. The antigen binding site is shaped by the variable domain of one heavy chain, along with that of its associated light chain, resulting in each immunoglobulin having</p>
-------------------	--

two antigen binding sites with remarkable affinity for a particular antigen. Variable domains undergo V-(D)-J rearrangement and subsequent somatic hypermutations, enabling affinity maturation for a specific antigen following exposure and selection. IgG1 protein mediates effector functions on monocytes, triggering antibody-dependent cellular cytotoxicity (ADCC) of virus-infected cells. Immunoglobulins are composed of two identical heavy chains and two identical light chains, interconnected by disulfide linkages, and interact with FCGR1A, FCGR2A, and FCGR3A to mediate various effector functions.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA