

OX40/TNFRSF4 Protein, Human (HEK 293, Fc)

Cat. No.:	HY-P70813
Synonyms:	Tumor necrosis factor receptor superfamily member 4; ACT35 antigen; OX40L receptor; TAX transcriptionally-activated glycoprotein 1 receptor; TNFRSF4; OX40; CD134; Txgp1
Species:	Human
Source:	HEK 293
Accession:	P43489 (L29-A216)
Gene ID:	7293
Molecular Weight:	Approximately 70.0 kDa

PROPERTIES

AA Sequence	<pre> L H C V G D T Y P S N D R C C H E C R P G N G M V S R C S R S Q N T V C R P C G P G F Y N D V V S S K P C K P C T W C N L R S G S E R K Q L C T A T Q D T V C R C R A G T Q P L D S Y K P G V D C A P C P P G H F S P G D N Q A C K P W T N C T L A G K H T L Q P A S N S S D A I C E D R D P P A T Q P Q E T Q G P P A R P I T V Q P T E A W P R T S Q G P S T R P V E V P G G R A V A </pre>
Biological Activity	2 µg/mL (100 µL/well) of immobilized Human OX40L-6His can bind Human OX40-Fc with an ED50 value of 4.23 µg/mL.
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O.
Storage & Stability	Stored at -20°C. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer. It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA