

BAFF Protein, Mouse (HEK 293, His)

Cat. No.:	HY-P70816
Synonyms:	Tumor necrosis factor ligand superfamily member 13B; B lymphocyte stimulator; BLyS; B-cell-activating factor; BAFF; Dendritic cell-derived TNF-like molecule; TNF- and APOL-related leukocyte expressed ligand 1; TALL-1
Species:	Mouse
Source:	HEK 293
Accession:	Q9WU72 (A127-L309)
Gene ID:	24099
Molecular Weight:	45-60 kDa

PROPERTIES

AA Sequence	<pre> A F Q G P E E T E Q D V D L S A P P A P C L P G C R H S Q H D D N G M N L R N I I Q D C L Q L I A D S D T P T I R K G T Y T F V P W L L S F K R G N A L E E K E N K I V V R Q T G Y F F I Y S Q V L Y T D P I F A M G H V I Q R K K V H V F G D E L S L V T L F R C I Q N M P K T L P N N S C Y S A G I A R L E E G D E I Q L A I P R E N A Q I S R N G D D T F F G A L K L L </pre>
Biological Activity	Human BCMA-His can bind Mouse BAFF-mFc with an affinity constant of <0.001 nM.
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O.
Storage & Stability	Stored at -20°C. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer. It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA