

TSLP Protein, Human (HEK293, His)

Cat. No.:	HY-P70818
Synonyms:	Thymic stromal lymphopoietin; TSLP
Species:	Human
Source:	HEK293
Accession:	Q969D9 (Y29-Q159)
Gene ID:	85480
Molecular Weight:	8-10&16-24 kDa

PROPERTIES

AA Sequence	<p>Y D F T N C D F E K I K A A Y L S T I S K D L I T Y M S G T K S T E F N N T V S</p> <p>C S N R P H C L T E I Q S L T F N P T A G C A S L A K E M F A M K T K A A L A I</p> <p>W C P G Y S E T Q I N A T Q A M K K R R K R K V T T N K C L E Q V S Q L Q G L W</p> <p>R R F N R P L L K Q Q</p>
Biological Activity	Immobilized Anti-Human TSLP mAb at 2 µg/mL (100 µl/well) can bind Human TSLP-His. The ED ₅₀ of Recombinant Human TSLP-His is <4.62 ng/mL.
Appearance	Lyophilized powder.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
Endotoxin Level	<1 EU/µg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH ₂ O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background	The TSLP protein is a cytokine that stimulates the release of T-cell-attracting chemokines from monocytes, with a particular effect on enhancing the maturation of CD11c(+) dendritic cells. Additionally, this protein can directly activate mast cells, potentially leading to the induction of allergic inflammation. It is also suggested that TSLP may have antimicrobial properties in the oral cavity and on the skin.
------------	---

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA