

## TRAIL R2/TNFRSF10B Protein, Mouse (HEK293, hFc)

<b>Cat. No.:</b>	HY-P70821
<b>Synonyms:</b>	Tumor necrosis factor receptor superfamily member 10B/Death receptor 5/MK/CD262/Tnfrsf10b/Dr5/Killer
<b>Species:</b>	Mouse
<b>Source:</b>	HEK293
<b>Accession:</b>	Q9QZM4 (N53-S177)
<b>Gene ID:</b>	21933
<b>Molecular Weight:</b>	50-75 kDa

### PROPERTIES

<b>AA Sequence</b>	<p>           N P A H N R P A G L    Q R P E E S P S R G    P C L A G Q Y L S E    G N C K P C R E G I            D Y T S H S N H S L    D S C I L C T V C K    E D K V V E T R C N    I T T N T V C R C K            P G T F E D K D S P    E I C Q S C S N C T    D G E E E L T S C T    P R E N R K C V S K            T A W A S         </p>
<b>Biological Activity</b>	Measured by its ability to inhibit TRAIL-mediated cytotoxicity using L-929 mouse fibroblast cells treated with TRAIL. The ED <sub>50</sub> this effect is 2-10 ng/mL in the presence of rhTRAIL, corresponding to a specific activity is 1-5×10 <sup>5</sup> units/mg.
<b>Appearance</b>	Lyophilized powder.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of 20 mM PB, 150 mM NaCl, pH 7.4.
<b>Endotoxin Level</b>	<1 EU/µg, determined by LAL method.
<b>Reconstitution</b>	It is not recommended to reconstitute to a concentration less than 100 µg/mL in ddH <sub>2</sub> O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).
<b>Storage &amp; Stability</b>	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
<b>Shipping</b>	Room temperature in continental US; may vary elsewhere.

### DESCRIPTION

<b>Background</b>	TRAIL R2/TNFRSF10B Protein serves as a receptor for the cytotoxic ligand TNFSF10/TRAIL. Upon ligand binding, the adapter molecule FADD recruits caspase-8 to the activated receptor, leading to the formation of the death-inducing signaling complex (DISC). The DISC performs caspase-8 proteolytic activation, initiating a cascade of caspases that mediate apoptosis. Additionally, TRAIL R2/TNFRSF10B promotes the activation of NF-κB and is essential for ER stress-induced apoptosis. In its monomeric form, it can interact with TRADD and RIPK1, and three TNFRSF10B molecules interact with the
-------------------	---

---

TNFSF10 homotrimer. In the absence of stimulation, TRAIL R2/TNFRSF10B interacts with BIRC2, DDX3X, and GSK3B, with enhanced interactions observed upon receptor stimulation, accompanied by DDX3X and BIRC2 cleavage (By similarity).

---

**Caution: Product has not been fully validated for medical applications. For research use only.**

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: [tech@MedChemExpress.com](mailto:tech@MedChemExpress.com)

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA