

TNF-alpha/TNFSF2 Protein, Mouse (P. pastoris)

Cat. No.:	HY-P7090A
Synonyms:	rMuTNF- α /TNFSF2; TNF-alpha; Cachectin; DIF; TNFA; Differentiation-inducing factor
Species:	Mouse
Source:	P. pastoris
Accession:	P06804 (S84-L235)
Gene ID:	21926
Molecular Weight:	Approximately 17 kDa

PROPERTIES

AA Sequence	S Q N S S D K P V A H V V A N H Q V E E Q L E W L S Q R A N A L L A N G M D L K D N Q L V V P A D G L Y L V Y S Q V L F K G Q G C P D Y V L L T H T V S R F A I S Y Q E K V N L L S A V K S P C P K D T P E G A E L K P W Y E P I Y L G G V F Q L E K G D Q L S A E V N L P K Y L D F A E S G Q V Y F G V I A L
Biological Activity	The ED ₅₀ is <0.01 ng/mL as measured by L929 cells, corresponding to a specific activity of >1 × 10 ⁸ units/mg.
Appearance	Lyophilized powder.
Formulation	Lyophilized after extensive dialysis against PBS.
Endotoxin Level	<1 EU/μg, determined by LAL method.
Reconstitution	It is not recommended to reconstitute to a concentration less than 100 μg/mL in ddH ₂ O. For long term storage it is recommended to add a carrier protein (0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose).
Storage & Stability	Stored at -20°C for 2 years. After reconstitution, it is stable at 4°C for 1 week or -20°C for longer (with carrier protein). It is recommended to freeze aliquots at -20°C or -80°C for extended storage.
Shipping	Room temperature in continental US; may vary elsewhere.

DESCRIPTION

Background	<p>TNF alpha is produced by various types of cells including macrophages, monocytes, neutrophils, T cells, and NK-cells^[2]. The amino acid sequence of human TNF alpha protein has low homology between mouse, rat, bovine, cynomolgus TNF alpha protein. While, human TNF alpha shares 94.85% aa sequence identity with cynomolgus TNF alpha protein, mouse TNF alpha shares 94.47% aa sequence identity with rat TNF alpha protein.</p> <p>TNF alpha exists in two forms; a type II transmembrane protein (tmTNF-α) and a mature soluble protein (sTNF-α). TNF-α binds to its receptors, mainly TNFR1 and TNFR2, and then transmits molecular signals for biological functions such as</p>
------------	---

inflammation and cell death. Both sTNF- α and tmTNF- α activate TNFR1, and process a death domain (DD) that interacts with the TNFR1-associated death domain (TRADD) adaptor protein. The TNFR2 signaling pathway is mainly activated by tmTNF- α . TNFR1 signaling tends to be pro-inflammatory and apoptotic. TNFR2 results in NF- κ B and MAPKs and AKT activation, TNFR2 activation is associated with homeostatic bioactivities such as tissue regeneration, cell proliferation, and cell survival, as well as host defense and inflammation^[1].

TNF-alpha is critical for normal immune response, abnormal secretion TNF alpha activates synovial fibroblasts, keratinocytes, osteoclasts, induces rheumatoid arthritis, inflammatory bowel disease, psoriatic arthritis (PsA), and noninfectious uveitis (NIU)^[3]. TNF alpha positively regulates endogenous TNF- α expression levels independently of Pgp efflux activity, induces IHF cells proliferation^[4]. TNF alpha in tissues may promote cancer growth, invasion, and metastasis. Besides, TNF alpha stimulates NF- κ B pathway via TNFR2 and anti-TNF- α MAb significantly suppresses the tumor development in colitis-associated cancer (CAC) mouse^[5]. TNF alpha as a proneurogenic factor activates the SAPK/JNK pathway and can facilitate neuronal replacement and brain repair in response to brain injury^[6].

REFERENCES

- [1]. Horiuchi T, et al. Transmembrane TNF-alpha: structure, function and interaction with anti-TNF agents. *Rheumatology (Oxford)*. 2010 Jul;49(7):1215-28.
- [2]. El-Tahan RR, et al. TNF- α gene polymorphisms and expression. *Springerplus*. 2016 Sep 7;5(1):1508.
- [3]. Jang DI, et al. The Role of Tumor Necrosis Factor Alpha (TNF- α) in Autoimmune Disease and Current TNF- α Inhibitors in Therapeutics. *Int J Mol Sci*. 2021 Mar 8;22(5):2719.
- [4]. Berguetti T, et al. TNF- α Modulates P-Glycoprotein Expression and Contributes to Cellular Proliferation via Extracellular Vesicles. *Cells*. 2019 May 24;8(5):500.
- [5]. Onizawa M, et al. Signaling pathway via TNF-alpha/NF-kappaB in intestinal epithelial cells may be directly involved in colitis-associated carcinogenesis. *Am J Physiol Gastrointest Liver Physiol*. 2009 Apr;296(4):G850-9.
- [6]. Bernardino L, et al. Tumor necrosis factor-alpha modulates survival, proliferation, and neuronal differentiation in neonatal subventricular zone cell cultures. *Stem Cells*. 2008 Sep;26(9):2361-71.
- [7]. Matsuno H, et al. The role of TNF-alpha in the pathogenesis of inflammation and joint destruction in rheumatoid arthritis (RA): a study using a human RA/SCID mouse chimera. *Rheumatology (Oxford)*. 2002 Mar;41(3):329-37.
- [8]. Zelová H, et al. TNF- α signalling and inflammation: interactions between old acquaintances. *Inflamm Res*. 2013 Jul;62(7):641-51.

Caution: Product has not been fully validated for medical applications. For research use only.

Tel: 609-228-6898

Fax: 609-228-5909

E-mail: tech@MedChemExpress.com

Address: 1 Deer Park Dr, Suite Q, Monmouth Junction, NJ 08852, USA